

DILATIONS OF SEMIGROUPS OF CONTRACTIONS THROUGH VESSELS

ELI SHAMOVICH AND VICTOR VINNIKOV

ABSTRACT. Let $A_1, \dots, A_d$ be a d -tuple of commuting dissipative operators on a separable Hilbert space $\mathcal{H}$. Using the theory of operator vessels and their associated systems, we give a construction of a dilation of the multi-parameter semigroup of contractions on $\mathcal{H}$ given by $e^{i \sum_{j=1}^d t_j A_j}$.

CONTENTS

1. Introduction	1
2. Livsic Commutative Operator Vessels	3
3. Solution in the Analytic Case	5
4. Very Reasonable Conditions	8
5. Solution in the Hyperbolic Case	12
6. Unitary Dilation of Semigroups	19
7. Construction of the Dilation (Proof of Theorem 6.4)	22
References	28

1. INTRODUCTION

The question of dilating a contraction to a unitary was considered first in 1953 by B. Sz.-Nagy in [40] (P. Halmos constructed a dilation in [21], but it was not a power dilation); the first explicit construction of a unitary dilation was given by J. Schäffer in [34]. In [40] Sz.-Nagy has also constructed dilations of one-parameter semigroups of contractions. Nagy's dilation theorem was extended to pairs of commuting contractions by T. Ando in [3]. However, in 1970 S. Parrot provided a counterexample to the existence of commuting unitary dilations for three commuting contractions in [29]. More examples appeared later, see for example [43]. In [8] and [6] W. Arveson generalized dilation theory to the setting of arbitrary operator algebras and their representations. More historical background and details are available in [7] and [41].

Our goal in this paper is to give an explicit construction of commutative unitary dilations of certain multi-parameter commutative semigroups of contractions. To be more precise, given a d -tuple of commuting dissipative operators on a separable Hilbert space $\mathcal{H}$, we consider the multiparameter semigroup S of contractions that they generate; we provide conditions on $A_1, \dots, A_d$, such that S admits a dilation to a commutative group of unitaries.

This paper is partially based on the results appearing in the Ph. D. thesis of E. S. written under the supervision of V. V. in the Ben-Gurion university of the Negev. Both authors were partially supported by US–Israel BSF grant 2010432.

As was already implicit in the works of M. S. Livsic (see [13]) and P. Lax and R. Phillips ([24]) and became explicit later ([1], [9], [22] and [23]), the construction of a unitary dilation has a simple system-theoretic interpretation: we embed the contraction into a conservative discrete-time input/state/output (i/s/o) linear system and consider the Hilbert space of square-summable trajectories with the natural shift operator. See also Sarason's Lemma [33, Lem. 0] that shows that any unitary dilation is obtained in this way, the works of B. Pavlov [31] and [30], and [10] for a survey of various multidimensional cases. We review these ideas in Section 7 (see Lemma 7.1 and Proposition 7.2). Motivated by this we construct an overdetermined multidimensional conservative (continuous-time) i/s/o system and consider a Hilbert space of certain trajectories of this system with a natural unitary representation of $\mathbb{R}^d$ on it by shifts. We expect that any commutative unitary dilation of a commutative semigroup of contractions arises in this way, so that the sufficient conditions on the operators $A_1, \dots, A_d$, that we describe, are necessary; we plan to address this question in a future work. For other sufficient conditions for the existence of a commutative unitary dilation in the discrete-time case, see [4], [5], [12], [16], [17] and [18].

In Section 2 we introduce the main tool in the construction, namely commutative operator vessels and their associated systems. More concretely, if $A_1, \dots, A_d$ is a d -tuple of commuting bounded operators on a separable Hilbert space $\mathcal{H}$, we associate to them a collection of spaces and operators:

$$\mathfrak{V} = (\mathcal{H}, \mathcal{E}, \Phi, \{A_j\}_{j=1}^d, \{\sigma_j\}_{j=1}^d, \{\gamma_{jk}\}_{j,k=1}^d, \{\gamma_{*jk}\}_{j,k=1}^d)$$

Here $\mathcal{E}$ is an auxiliary separable Hilbert space, $\Phi: \mathcal{H} \rightarrow \mathcal{E}$ is a bounded operator and for every $j, k = 1, \dots, d$, σ_j , γ_{jk} and γ_{*jk} are selfadjoint bounded operators on $\mathcal{E}$ satisfying some conditions described in detail in the text (see (2.1) and (2.2)). The study of operator vessels was initiated by M. S. Livsic (see for example the papers [13], [28], [25] and [26] and the book [27]). A functional model for two commuting dissipative operators with finite-dimensional imaginary parts was constructed by J. Ball and the second author in [11] using frequency domain methods. We briefly recall the relevant notions of the associated overdetermined multidimensional system, the adjoint system and the input and output compatibility conditions in both the continuous and the discrete-time setting. We introduce all the necessary background, notions and results. Some of the results are proved for the sake of completeness. In particular we show that there is a natural way, given a d -tuple of commuting operators, to embed them in a so-called strict vessel.

In Section 3 we consider the system of input and output compatibility conditions in the analytic case. If $d \geq 3$ this system is itself overdetermined. We find therefore necessary and sufficient conditions on the vessel, so that the system of input (or output) compatibility conditions admits a solution for any initial condition along one of the axes analytic in some neighborhood of the origin. We call these conditions very reasonable conditions or VR for short. We then show in Section 4 that the VR conditions are independent of the choice of the axis and they hold at the input if and only if they hold at the output. We also show that in the case of a doubly commuting d -tuple of operators, we have the VR conditions automatically for the strict vessel embedding.

We proceed in Section 5 to show that if the VR conditions hold and the system of continuous-time compatibility conditions is hyperbolic, then it has a weak solution in tempered distributions for every initial condition along one of the axes. We

proceed to show that under certain assumptions the distribution is in fact a function that is in L^2 on lines with respect to a twisted inner product. We write a transform taking the initial condition along the t_1 axis into a condition along the t_j axis and demonstrate some of its properties. This section forms the technical toolbox for the proof of the main dilation theorem.

In Section 6 we state the main dilation theorem: if a d -tuple of commuting dissipative operators $A_1, \dots, A_d$ possesses the dissipative embedding property, namely if they can be embedded into a vessel satisfying the VR conditions and such that $\sigma_1, \dots, \sigma_d \geq 0$, then the semigroup of contraction generated by $A_1, \dots, A_d$ admits a dilation to a commutative group of unitaries. In particular since the VR conditions are vacuous when $d = 2$ we obtain a version of Ando's dilation theorem for the continuous-time case. There are some technical restrictions, since we are dealing with bounded generators, however our construction of the unitary dilation is completely explicit and thus should allow a further geometric analysis, similarly to the one dimensional case.

We prove the main dilation theorem, i.e., we construct the dilation space and the group of unitaries in Section 7 using the tools developed in Section 5. We conclude by demonstrating a necessary and a sufficient condition for the dilation thus obtained to be minimal.

2. LIVSIC COMMUTATIVE OPERATOR VESSELS

In this section we recall briefly the notion of Livsic commutative operator vessels, for more information see [27].

Definition 2.1. Let $\mathcal{H}$ be a separable Hilbert space and $A_1, \dots, A_d$ a d -tuple of commuting non-selfadjoint bounded operators on $\mathcal{H}$. We fix an auxiliary separable Hilbert space $\mathcal{E}$, a bounded operator $\Phi: \mathcal{H} \rightarrow \mathcal{E}$ and a d -tuple of bounded selfadjoint operators on $\mathcal{E}$, $\sigma_1, \dots, \sigma_d$, satisfying the colligation condition, namely for every $k = 1, \dots, d$:

$$(2.1) \quad A_k - A_k^* = i\Phi^* \sigma_k \Phi.$$

We also fix two collections of bounded selfadjoint operators, γ_{jk} and γ_{*jk} , for $j, k = 1, \dots, d$ on $\mathcal{E}$, satisfying the following set of conditions:

$$(2.2) \quad \begin{aligned} &\bullet \gamma_{jk} = -\gamma_{kj}, \gamma_{*jk} = -\gamma_{*kj}, \\ &\bullet \sigma_j \Phi A_k^* - \sigma_k \Phi A_j^* = \gamma_{jk} \Phi, \\ &\bullet \sigma_j \Phi A_k - \sigma_k \Phi A_j = \gamma_{*jk} \Phi, \\ &\bullet \gamma_{*jk} - \gamma_{jk} = i(\sigma_j \Phi \Phi^* \sigma_k - \sigma_k \Phi \Phi^* \sigma_j). \end{aligned}$$

The collection of operators and spaces satisfying the above conditions is called a Livsic commutative operator vessel.

Given a Livsic commutative vessel one can associate to it an energy preserving linear time invariant overdetermined system in continuous-time. Let $u, y: \mathbb{R}^d \rightarrow \mathcal{E}$ be smooth functions, we call them the input and output signals, respectively, and let $x: \mathbb{R}^d \rightarrow \mathcal{H}$ be a smooth function that we call the state, then we define the system:

$$\begin{aligned} i \frac{\partial x}{\partial t_k} + A_k x &= \Phi^* \sigma_k u, \\ y &= u - i\Phi x. \end{aligned}$$

We will assume for now that $u, y \in C^1(\mathbb{R}^d, \mathcal{E})$ and $x \in C^2(\mathbb{R}^d, \mathcal{H})$. We will discuss latter on various relaxations of this assumption.

For $d > 1$ the above system is overdetermined and hence requires input and output compatibility conditions. It follows from the vessel conditions (2.2) (for details see [27, Thm. 3.2.1]) that the necessary and sufficient input compatibility conditions are given by:

$$(2.3) \quad \Phi^* \left(\sigma_k \frac{\partial u}{\partial t_j} - \sigma_j \frac{\partial u}{\partial t_k} + i\gamma_{jk} u \right) = 0.$$

We define the strict input compatibility conditions by:

$$(2.4) \quad \sigma_k \frac{\partial u}{\partial t_j} - \sigma_j \frac{\partial u}{\partial t_k} + i\gamma_{jk} u = 0.$$

Similarly at the output we get the following system of compatibility conditions and strict compatibility conditions:

$$(2.5) \quad \Phi^* \left(\sigma_k \frac{\partial y}{\partial t_j} - \sigma_j \frac{\partial y}{\partial t_k} + i\gamma_{*jk} y \right) = 0.$$

$$(2.6) \quad \sigma_k \frac{\partial y}{\partial t_j} - \sigma_j \frac{\partial y}{\partial t_k} + i\gamma_{*jk} y = 0.$$

When $d > 2$ we note that the system of input compatibility conditions (2.4) is itself overdetermined. The goal of the current paper is to understand the additional compatibility conditions on (2.4) required for the system to have "enough" solutions in the hyperbolic case, i.e., when the operators $A_1, \dots, A_d$ are dissipative and $\sigma_1, \dots, \sigma_d \geq 0$. We then use these solutions to construct a unitary dilation for the semigroup of contractions generated by $A_1, \dots, A_d$.

One can show that if u solves the system of input compatibility conditions, then for each initial condition $x(0) = h \in \mathcal{H}$, there exists a unique state x solving the system and the output y then satisfies the output compatibility conditions, see [11] for the $d = 2$ case and [27, 36] for the general case. The formula for x is then:

$$(2.7) \quad x(t_1, \dots, t_d) = e^{i \sum_{j=1}^d t_j A_j} (h - i \int_0^{(t_1, \dots, t_d)} e^{-i \sum_{j=1}^d s_j A_j} \Phi^* \left(\sum_{j=1}^d \sigma_j u(s_1, \dots, s_d) ds_j \right) ds).$$

We also have the adjoint vessel:

$$\mathfrak{V}^* = (\mathcal{H}, \mathcal{E}, -\Phi, \{A_j^*\}_{j=1}^d, \{-\sigma_j\}_{j=1}^d, \{-\gamma_{jk}\}_{j,k=1}^d, \{-\gamma_{*jk}\}_{j,k=1}^d)$$

The adjoint system, namely the associated system of the adjoint vessel, is given by:

$$(2.8) \quad \begin{aligned} i \frac{\partial \tilde{x}}{\partial t_k} + A_k^* \tilde{x} &= \Phi^* \sigma_k \tilde{u}, \\ \tilde{y} &= \tilde{u} + i\Phi \tilde{x}. \end{aligned}$$

It is proved in [11], [27] and [36] that (u, x, y) is a system trajectory for the associated system of $\mathfrak{V}$ if and only if (y, x, u) is a system trajectory for the adjoint system. Using this we deduce the energy balance equations ([36, Cor. 1.2.8]).

Namely for a trajectory (u, x, y) of the associated system we have for every $t = (t_1, \dots, t_d) \in \mathbb{R}^d$:

$$(2.9) \quad \|x(t + se_j)\|^2 - \|x(t)\|^2 = \int_0^s \langle \sigma_j u(t + pe_j), u(t + pe_j) \rangle dp - \int_0^s \langle \sigma_j y(t + pe_j), y(t + pe_j) \rangle dp.$$

We briefly recall the proof for the sake of completeness:

$$\begin{aligned} \frac{\partial}{\partial t_j} \langle x, x \rangle &= \left\langle \frac{\partial x}{\partial t_j}, x \right\rangle + \left\langle x, \frac{\partial x}{\partial t_j} \right\rangle = \langle iA_j x - i\Phi^* \sigma_j u, x \rangle + \langle x, iA_j x - i\Phi^* \sigma_j u \rangle = \\ &= \langle i(A_j - A_j^*)x, x \rangle - i\langle u, \sigma_j \Phi x \rangle + i\langle \sigma_j \Phi x, u \rangle = \\ &= \langle u, \sigma_j (u - y) \rangle + \langle \sigma_j (u - y), u \rangle - \langle \sigma_j \Phi x, \Phi x \rangle = \langle \sigma_j u, u \rangle - \langle \sigma_j y, y \rangle. \end{aligned}$$

Now all that remains is to integrate with respect to t_j to get the desired result. Note that it follows from this proof that if (u, x, y) is a system trajectory, with u and y locally integrable on every line parallel to one of the axes and such that the system still admits a solution x that is absolutely continuous on every such line and thus almost everywhere differentiable on it (as a function of one variable), then the energy balance equations still hold. We will use this comment in the following sections.

Definition 2.2. We will say that a vessel $\mathfrak{V}$ is strict if:

- Φ is surjective,
- $\cap_{j=1}^d \ker \sigma_j = 0$.

We will say that $\mathfrak{V}$ is weakly strict if $\cap_{j=1}^d \ker \Phi^* \sigma_j = 0$.

Clearly, if $\mathfrak{V}$ is strict it is weakly strict; the converse is not necessarily true. It was shown in [27] that every d -tuple of commuting operators admits an embedding into an essentially unique strict vessel (see [36] for a non-commutative case). The embedding is given as follows:

$$(2.10) \quad \begin{aligned} \mathcal{E} &= \overline{\sum_{j=1}^d \text{Im}(A_j - A_j^*)}, \\ \Phi &= P_{\mathcal{E}}, \\ \sigma_j &= \frac{1}{i}(A_j - A_j^*)|_{\mathcal{E}}, \\ \gamma_{jk} &= \frac{1}{i}(A_j A_k^* - A_k A_j^*)|_{\mathcal{E}}, \\ \gamma_{*j k} &= \frac{1}{i}(A_j^* A_k - A_k^* A_j)|_{\mathcal{E}}. \end{aligned}$$

The subspace $\mathcal{E}$ is called the non-Hermitian subspace of the d -tuple $A_1, \dots, A_d$.

3. SOLUTION IN THE ANALYTIC CASE

Assume that u is a real analytic function with a convergent power series expansion around the origin:

$$u(t) = \sum_{n \in \mathbb{N}^d} a(n) t^n.$$

Then plugging the power series into (2.4) we get the following difference equation on the coefficients of the power series:

$$\sigma_k a(n + e_j) - \sigma_j a(n + e_k) + i\gamma_{jk} a(n) = 0.$$

Remark 3.1. Consider a collection of operators and spaces satisfying the vessel conditions (2.2), i.e., a vessel in the more general sense of [11] and [27, Part III] (without the colligation condition (2.1)). We can associate to such a general vessel a linear overdetermined discrete-time i/s/o system, such that the corresponding system of input compatibility conditions is the system of difference equations obtained above.

We will assume from now on that σ_1 is invertible. Denote by d_j the shift operator in the j -th coordinate, namely $(d_j a)(n) = a(n + e_j)$, for every $n \in \mathbb{N}^d$. We would like to be able to solve the system for every initial condition along the n_1 axis. Consider an equation that contains σ_1 , and multiply it by σ_1^{-1} to get:

$$(3.1) \quad a(n + e_k) = \sigma_1^{-1} \sigma_k a(n + e_1) + i\sigma_1^{-1} \gamma_{1k} a(n).$$

Applying first d_j and then d_k we get:

$$\begin{aligned} a(n + e_k + e_j) &= \sigma_1^{-1} \sigma_k a(n + e_j + e_1) + i\sigma_1^{-1} \gamma_{1k} a(n + e_j) = \\ &= \sigma_1^{-1} \sigma_k \sigma_1^{-1} \sigma_j a(n + 2e_1) + i\sigma_1^{-1} \sigma_k \sigma_1^{-1} \gamma_{1j} a(n + e_1) + i\sigma_1^{-1} \gamma_{1k} \sigma_1^{-1} \sigma_j a(n + e_1) - \\ &\quad \sigma_1^{-1} \gamma_{1k} \sigma_1^{-1} \gamma_{1j} a(n). \end{aligned}$$

Since the shift operators along different axes commute we get that:

$$\begin{aligned} &(\sigma_k \sigma_1^{-1} \sigma_j - \sigma_k \sigma_1^{-1} \sigma_j) a(n + 2e_1) + (\gamma_{1j} \sigma_1^{-1} \gamma_{1k} - \gamma_{1k} \sigma_1^{-1} \gamma_{1j}) a(n) + \\ &\quad i(\sigma_k \sigma_1^{-1} \gamma_{1j} + \gamma_{1k} \sigma_1^{-1} \sigma_j - \sigma_j \sigma_1^{-1} \gamma_{1k} - \gamma_{1j} \sigma_1^{-1} \sigma_k) a(n + e_1) = 0. \end{aligned}$$

Now if we take $n = 0$ and use the fact that we require the system to be solvable for every initial condition along the n_1 -axis we get the following set of necessary conditions:

$$(3.2) \quad \begin{aligned} &\bullet [\sigma_1^{-1} \sigma_j, \sigma_1^{-1} \sigma_k] = 0, \\ &\bullet [\sigma_1^{-1} \gamma_{1j}, \sigma_1^{-1} \gamma_{1k}] = 0, \\ &\bullet [\sigma_1^{-1} \sigma_k, \sigma_1^{-1} \gamma_{1j}] = [\sigma_1^{-1} \sigma_j, \sigma_1^{-1} \gamma_{1k}]. \end{aligned}$$

There are more necessary conditions, since we have a lot of equations that do not involve σ_1 . We take such an equation and use (3.1) to get:

$$\begin{aligned} &\sigma_k \sigma_1^{-1} \sigma_j a(n + e_1) + i\sigma_k \sigma_1^{-1} \gamma_{1j} a(n) - \sigma_j \sigma_1^{-1} \sigma_k a(n + e_1) - \\ &\quad i\sigma_j \sigma_1^{-1} \gamma_{1k} a(n) + i\gamma_{jk} a(n) = 0. \end{aligned}$$

If we apply now the necessary conditions (3.2) and use again the fact that for $n = 0$ the vector $a(0)$ is arbitrary we get that:

$$(3.3) \quad \gamma_{jk} = \sigma_j \sigma_1^{-1} \gamma_{1k} - \sigma_k \sigma_1^{-1} \gamma_{1j}.$$

Then the following proposition is almost immediate:

Proposition 3.2. *The conditions (3.2) and (3.3) are necessary and sufficient for the system of discrete-time input compatibility equations to have a solution for every initial condition along the n_1 -axis.*

Proof. We have seen above that this set of conditions is necessary. Now for sufficiency note that using (3.3) we can eliminate all of the equations that do not involve σ_1 . The other equations are compatible by (3.2) and thus for every initial $a(n_1, 0, \dots, 0) = b(n_1)$ they define a unique function a . In fact if $a(n_1, 0, \dots, 0) = b(n_1)$ we can write a as follows (recall that d_1 stands for the shift in the first coordinate):

$$(3.4) \quad a(n_1, \dots, n_d) = (\alpha_2 d_1 + i\beta_2)^{n_2} \cdots (\alpha_d d_1 + i\beta_d)^{n_d} b(n_1).$$

Here $\alpha_j = \sigma_1^{-1} \sigma_j$ and $\beta_j = \sigma_1^{-1} \gamma_{1j}$. □

Remark 3.3. For the necessity part of Proposition 3.2 it is in fact enough to assume that $a(0)$, $a(e_1)$ and $a(2e_1)$ are arbitrary.

Definition 3.4. We will call a vessel that satisfies conditions (3.2) and (3.3) very reasonable.

Remark 3.5. In [37, Cor. 2.20] and the following discussion, similar conditions were given for a tensor $\gamma \in M_n(\mathbb{C}) \otimes \wedge^{k+1} \mathbb{C}^{d+1}$ to be very reasonable. The difference is that in [37] we require $\mathcal{E}$ to be finite-dimensional and we also require generic semisimplicity, whereas in the case at hand we do not need either.

We now use (3.2) and (3.3) to describe the solution in the continuous-time case when the initial condition is an $\mathcal{E}$ -valued analytic function in a neighbourhood of 0. Recall, that a function $f: (-r, r) \rightarrow \mathcal{E}$ is strongly analytic at 0 if there exist $\{\xi_n\}_{n=0}^\infty \subset \mathcal{E}$, such that $f(t) = \sum_{n=0}^\infty \xi_n t^n$, where the series converges in norm for every t in a neighborhood of 0. This condition is in fact equivalent to weak analyticity, namely that for every $\xi \in \mathcal{E}$ the function $\langle f(t), \xi \rangle$ is real analytic in a neighbourhood of 0. The following theorem can be thought of as a version of the classical Cauchy-Kowalevskaya theorem (cf. [15, Sec. I.D]).

Theorem 3.6. Assume that we are given an initial condition $u(t_1, 0, \dots, 0) = f(t_1)$ analytic near the origin and that (3.2) and (3.3) are satisfied, then there exists an open neighborhood of the origin and a unique analytic solution u to the input compatibility system.

Proof. First note that as in the discrete-time case, (3.3) allows us to eliminate all of the equations that do not involve σ_1 , hence we are left with the system ($j = 2, \dots, d$):

$$\frac{\partial u}{\partial t_j} = \alpha_j \frac{\partial u}{\partial t_1} + \beta_j u.$$

Here $\alpha_j = \sigma_1^{-1} \sigma_j$ and $\beta_j = i\sigma_1^{-1} \gamma_{1j}$. We write an expansion for the initial condition f and the solution u in a polydisc around the origin and solve for the coefficients. This reduces the problem to the discrete-time case that we have already seen, except that we have to verify that the series for u is locally convergent. If the radius of convergence of the series of $f(t_1) = \sum_{m=0}^\infty b(m)t_1^m$ around 0 is R , then for every $0 < r < R$, there exists a constant $M > 0$, such that $\|b(m)\| \leq \frac{M}{r^m}$. Now we note that we can obtain the coefficients of u in terms of the $b(m)$ using Equation 3.4. If we set $C = \max\{\|\alpha_j\|, \|\beta_j\| \mid j = 2, \dots, d\}$, then we get that:

$$\|a(n)\| \leq C^{|n|-n_1} M \frac{1}{r^{n_1}} \left(1 + \frac{1}{r}\right)^{|n|-n_1}.$$

Therefore, the series for u will converge in the closed polydisc around the origin with polyradius $(r, \frac{r}{C(r+1)}, \dots, \frac{r}{C(r+1)})$. Since this is true for every $0 < r < R$, we see that the series for u will converge in the polydisc around the origin with polyradius $(R, \frac{R}{C(R+1)}, \dots, \frac{R}{C(R+1)})$. $\square$

4. VERY REASONABLE CONDITIONS

In this Section we will discuss the conditions (3.2) and (3.3) that we will call very reasonable conditions or VR conditions for brevity. We have defined the VR conditions in the case of the input compatibility system. A similar set of conditions arises at the output, namely for $j, k = 2, \dots, d$:

$$(4.1) \quad \begin{aligned} & \bullet [\sigma_1^{-1}\sigma_j, \sigma_1^{-1}\sigma_k] = 0, \\ & \bullet [\sigma_1^{-1}\gamma_{*1j}, \sigma_1^{-1}\gamma_{*1k}] = 0, \\ & \bullet [\sigma_1^{-1}\sigma_k, \sigma_1^{-1}\gamma_{*1j}] = [\sigma_1^{-1}\sigma_j, \sigma_1^{-1}\gamma_{*1k}], \\ & \bullet \gamma_{*jk} = \sigma_j\sigma_1^{-1}\gamma_{*1k} - \sigma_k\sigma_1^{-1}\gamma_{*1j}. \end{aligned}$$

Let us call this system of conditions VR_* conditions. We will now investigate the relation between the VR and VR_* conditions.

Proposition 4.1. *Given a vessel $\mathfrak{V}$, it satisfies the VR conditions if and only if it satisfies the VR_* conditions.*

Proof. It suffices to prove only one implication, since the proof of the other will be symmetric. Assume that the VR conditions hold and thus the first condition of the VR_* conditions is automatically satisfied.

From the linkage vessel condition, we get:

$$(4.2) \quad \gamma_{jk} = \gamma_{*jk} - i\sigma_j\Phi\Phi^*\sigma_k + i\sigma_k\Phi\Phi^*\sigma_j.$$

To prove that the fourth compatibility condition of (4.1) holds, we use the linkage condition to obtain:

$$\gamma_{*jk} = \gamma_{jk} + i\sigma_j\Phi\Phi^*\sigma_k - i\sigma_k\Phi\Phi^*\sigma_j.$$

Then equation (3.3) yields:

$$(4.3) \quad \gamma_{*jk} = \sigma_j\sigma_1^{-1}\gamma_{1k} - \sigma_k\sigma_1^{-1}\gamma_{1j} + i\sigma_j\Phi\Phi^*\sigma_k - i\sigma_k\Phi\Phi^*\sigma_j.$$

Now using (4.2), we get:

$$(4.4) \quad \begin{aligned} \sigma_j\sigma_1^{-1}\gamma_{1k} &= \sigma_j\sigma_1^{-1}\gamma_{*1k} - i\sigma_j\Phi\Phi^*\sigma_k + i\sigma_j\sigma_1^{-1}\sigma_k\Phi\Phi^*\sigma_1 \\ \gamma_{1k}\sigma_1^{-1}\sigma_j &= \sigma_j\sigma_1^{-1}\gamma_{*1k} - i\sigma_1\Phi\Phi^*\sigma_k\sigma_1^{-1}\sigma_j + i\sigma_k\Phi\Phi^*\sigma_j \end{aligned}$$

Plugging in the equations of (4.4) into (4.3), and using the first condition of (3.2), we obtain the fourth equation of (4.1).

Similarly we consider the third equation of (3.2). We use (4.4) and obtain immediately the third condition of (4.1).

Using the vessel conditions one obtains the following equation:

$$(4.5) \quad \begin{aligned} \sigma_j\Phi\Phi^*\sigma_k\Phi\Phi^*\sigma_l &= i\sigma_j\Phi A_k^*\Phi^*\sigma_k - i\sigma_j\Phi A_k\Phi^*\sigma_l \\ i\gamma_{*jk}\Phi\Phi^*\sigma_l &= i\sigma_k\Phi A_j\Phi^*\sigma_l - i\sigma_j\Phi A_k\Phi^*\sigma_l \\ i\sigma_j\Phi\Phi^*\gamma_{*kl} &= i\sigma_j\Phi A_k^*\Phi^*\sigma_l - i\sigma_j\Phi A_l^*\Phi^*\sigma_k. \end{aligned}$$

Using (4.4) on the second condition of (3.2). we get:

$$\begin{aligned}
(4.6) \quad 0 &= \gamma_{*1j}\sigma_1^{-1}\gamma_{*1k} - \gamma_{*1k}\sigma_1^{-1}\gamma_{*1j} - i\gamma_{*1j}\Phi\Phi^*\sigma_k + \\
&+ i\gamma_{*1k}\Phi\Phi^*\sigma_j + i(\gamma_{*1j}\sigma_1^{-1}\sigma_k - \gamma_{*1k}\sigma_1^{-1}\sigma_j)\Phi\Phi^*\sigma_1 + \\
&+ i\sigma_j\Phi\Phi^*\gamma_{*1k} - i\sigma_3\Phi\Phi^*\gamma_{*1j} + i\sigma_1\Phi\Phi^*(\sigma_k\sigma_1^{-1}\gamma_{*1j} - \sigma_j\sigma_1^{-1}\gamma_{*1k}) - \\
&- \sigma_1\Phi\Phi^*\sigma_j\Phi\Phi^*\sigma_k + \sigma_j\Phi\Phi^*\sigma_1\Phi\Phi^*\sigma_k - \sigma_j\Phi\Phi^*\sigma_j\Phi\Phi^*\sigma_1 + \\
&+ \sigma_1\Phi\Phi^*\sigma_j\Phi\Phi^*\sigma_k - \sigma_k\Phi\Phi^*\sigma_1\Phi\Phi^*\sigma_j + \sigma_k\Phi\Phi^*\sigma_j\Phi\Phi^*\sigma_1
\end{aligned}$$

Now using the fourth equation of (4.1) and (4.5), we see that all the terms cancel, but for the first two. Thus we have obtained the second equation of (4.1) and the proposition is proved. $\square$

Now that we know that the VR conditions fit naturally into the framework of vessels, we ask a question about invariance under coordinate changes. Namely, we assume that σ_2 is invertible as well and we can write a system of VR conditions for σ_2 , for $j, k = 1, 3, 4, \dots, d$:

$$\begin{aligned}
(4.7) \quad &\bullet [\sigma_2^{-1}\sigma_j, \sigma_2^{-1}\sigma_k] = 0, \\
&\bullet [\sigma_2^{-1}\gamma_{2j}, \sigma_2^{-1}\gamma_{2k}] = 0, \\
&\bullet [\sigma_2^{-1}\sigma_k, \sigma_2^{-1}\gamma_{2j}] = [\sigma_2^{-1}\sigma_j, \sigma_2^{-1}\gamma_{2k}], \\
&\bullet \gamma_{jk} = \sigma_j\sigma_2^{-1}\gamma_{2k} - \sigma_k\sigma_2^{-1}\gamma_{2j}.
\end{aligned}$$

We will call this system of conditions VR conditions in the direction of e_2 and we refer to the original VR conditions as VR conditions in the direction of e_1 .

Proposition 4.2. *Given a vessel $\mathfrak{V}$ such that both σ_1 and σ_2 are invertible, then the VR conditions in the direction of e_1 are satisfied if and only if the VR conditions in the direction of e_2 are satisfied.*

Proof. Let us assume that the VR conditions in the direction of e_2 hold. Then for every $k = 3, \dots, d$ we have that:

$$\sigma_1\sigma_2^{-1}\sigma_k = \sigma_k\sigma_2^{-1}\sigma_1.$$

Premultiplying by σ_1^{-1} we get:

$$\sigma_2^{-1}\sigma_k = \sigma_1^{-1}\sigma_k\sigma_2^{-1}\sigma_1.$$

Hence for every $j, k = 3, \dots, d$ we get:

$$\sigma_j\sigma_2^{-1}\sigma_k = \sigma_j\sigma_1^{-1}\sigma_k\sigma_2^{-1}\sigma_1.$$

Now using the first VR condition for σ_2 we get:

$$\sigma_j\sigma_1^{-1}\sigma_k\sigma_2^{-1}\sigma_1 = \sigma_k\sigma_1^{-1}\sigma_j\sigma_2^{-1}\sigma_1.$$

This gives us the first VR condition in the direction of e_1 , namely the first equation of (3.2) for $j, k = 3, \dots, d$. We only need to check for $j = 2$ and $k = 3, \dots, d$:

$$\begin{aligned}
\sigma_2\sigma_1^{-1}\sigma_k - \sigma_k\sigma_1^{-1}\sigma_2 &= \sigma_2\sigma_1^{-1}(\sigma_k - \sigma_1\sigma_2^{-1}\sigma_k\sigma_1^{-1}\sigma_2) = \\
&= \sigma_2\sigma_1^{-1}(\sigma_k - \sigma_k\sigma_2^{-1}\sigma_1\sigma_1^{-1}\sigma_2) = 0.
\end{aligned}$$

We get the second equation of (3.2) in exactly the same way. Combining the two we get easily the third equation of (3.2).

Now we use the fourth equation of (4.7) to get that for every $k = 3, \dots, d$:

$$\gamma_{1k} = \sigma_1 \sigma_2^{-1} \gamma_{2k} - \sigma_k \sigma_2^{-1} \gamma_{21}.$$

Therefore for $j, k = 2, \dots, d$:

$$\begin{aligned} \sigma_j \sigma_1^{-1} \gamma_{1k} - \sigma_k \sigma_1^{-1} \gamma_{1j} &= \sigma_j \sigma_2^{-1} \gamma_{2k} - \sigma_j \sigma_1^{-1} \sigma_k \sigma_2^{-1} \gamma_{21} \\ &\quad - \sigma_k \sigma_2^{-1} \gamma_{2j} + \sigma_k \sigma_1^{-1} \sigma_j \sigma_2^{-1} \gamma_{21} = \gamma_{jk}. \end{aligned}$$

Here we have used the first equation of (3.2) and the fourth equation of (4.7) for $j, k = 3, \dots, d$. For $j = 2$ and $k = 3, \dots, d$ we get:

$$\begin{aligned} \sigma_2 \sigma_1^{-1} \gamma_{1k} - \sigma_k \sigma_1^{-1} \gamma_{12} &= \\ \gamma_{2k} - \sigma_2 \sigma_1^{-1} \sigma_k \sigma_2^{-1} \gamma_{21} - \sigma_k \sigma_1^{-1} \gamma_{12} &= \gamma_{2k}. \end{aligned}$$

Hence we proved that (3.3) holds. $\square$

Given a vessel $\mathfrak{V}$ we can consider as in [36] the linear maps $\rho: \mathbb{R}^d \rightarrow B(\mathcal{H})$, $\sigma: \mathbb{R}^d \rightarrow B(\mathcal{H})$ and $\gamma: \wedge^2 \mathbb{R}^d \rightarrow B(\mathcal{H})$ given by:

$$\rho(e_j) = A_j, \sigma(e_j) = \sigma_j, \gamma(e_j \wedge e_k) = \gamma_{jk}.$$

Then for every $T \in \text{GL}_d(\mathbb{R})$ we can define:

$$A_j^T = \rho(Te_j), \sigma_j^T = \sigma(Te_j), \gamma_{jk}^T = \gamma(\wedge^2(T)e_j \wedge e_k).$$

Thus we get a vessel $\mathfrak{V}^T$, we call this the coordinate change corresponding to T . We will say that $\mathfrak{V}$ satisfies the VR conditions in the direction of Te_1 if $\mathfrak{V}^T$ satisfies the VR conditions in the direction of e_1 , generalizing (4.7).

Corollary 4.3. *Let us assume that there exist $\xi, \eta \in \mathbb{R}^d$, such that both $\xi\sigma = \sum_{j=1}^d \xi_j \sigma_j$ and $\eta\sigma$ are invertible and $\mathfrak{V}$ satisfies the VR conditions in the direction of ξ then it satisfies the VR conditions in the direction of η . In particular, there exists an open set $U \subset \text{GL}_d(\mathbb{R})$, such that for every $T \in U$, $\mathfrak{V}$ satisfies the VR conditions in the direction of $T\xi$.*

Proof. We can take $T \in \text{GL}_d(\mathbb{R})$, such that $T\xi = e_1$ and $T\eta = e_2$ and apply Proposition 4.2. To obtain the second part of the statement we note that since the invertible matrices are an open set, for $\eta \in \mathbb{R}^d$, such that $\|\xi\sigma - \eta\sigma\|$ is small enough, we have that $\eta\sigma$ is invertible. $\square$

This corollary allows us to treat VR conditions without mentioning the direction. For definiteness we will assume for the rest of this section that the VR conditions in direction e_1 are satisfied.

Remark 4.4. In case $\dim \mathcal{E} < \infty$ one notes that the set U from the above Corollary is in fact Zariski open and dense.

The VR conditions are slightly redundant as the following proposition shows:

Proposition 4.5. *Assume that $\mathfrak{V}$ is a vessel that satisfies (3.3), then it satisfies the third condition of (3.2) automatically.*

Proof. Since γ_{jk} and σ_j are selfadjoint we get by taking the adjoint of (3.3) that:

$$\gamma_{jk} = \gamma_{1k} \sigma_1^{-1} \sigma_j - \gamma_{1j} \sigma_1^{-1} \sigma_k.$$

Now subtract it from (3.3) to get the third equation of (3.2). $\square$

The following is a strong converse to Proposition 4.5 and provides a way to construct VR vessels from partial data:

Proposition 4.6. *Assume that we are given a d -tuple of commuting non-selfadjoint operators $A_1, \dots, A_d$ on $\mathcal{H}$, an operator $\Phi: \mathcal{H} \rightarrow \mathcal{E}$ and a collection of selfadjoint operators $\sigma_1, \dots, \sigma_d$ and $\gamma_{12}, \dots, \gamma_{1d}$ on $\mathcal{E}$, such that σ_1 is invertible, the commutativity conditions (3.2) hold and the relevant vessel conditions hold, namely for every $j = 1, \dots, d$:*

$$\begin{aligned} A_j - A_j^* &= i\Phi^* \sigma_j \Phi. \\ \gamma_{1j} \Phi &= \sigma_1 \Phi A_j^* - \sigma_j \Phi A_1^*. \end{aligned}$$

Then there exists a VR vessel $\mathfrak{V}$ with the above data.

Proof. We define γ_{jk} using (3.3), namely:

$$\gamma_{jk} = \sigma_j \sigma_1^{-1} \gamma_{1k} - \sigma_k \sigma_1^{-1} \gamma_{1j}.$$

Then using the same computation as in the preceding Proposition we see that γ_{jk} is selfadjoint. Now to see that it satisfies the input vessel condition we check:

$$\begin{aligned} \gamma_{jk} \Phi &= \sigma_j \sigma_1^{-1} \gamma_{1k} \Phi - \sigma_k \sigma_1^{-1} \gamma_{1j} \Phi = \\ &= \sigma_j \sigma_1^{-1} (\sigma_1 \Phi A_k^* - \sigma_k \Phi A_1^*) - \sigma_k \sigma_1^{-1} (\sigma_1 \Phi A_j^* - \sigma_j \Phi A_1^*) = \sigma_j \Phi A_k^* - \sigma_k \Phi A_j^*. \end{aligned}$$

The last equality follows from the first condition of (3.2). Now we define γ_{*jk} using the linkage condition to get a vessel $\mathfrak{V}$. It is obvious that this vessel satisfies the VR conditions. $\square$

Recall that from Definition 2.2 a strict vessel is a vessel, such that Φ is surjective and $\cap_{j=1}^d \ker \sigma_j = 0$. Since we assume that σ_1 is invertible, the second condition holds automatically. The strict vessels are slightly easier to work with as the following claim shows:

Proposition 4.7. *Assume that $\mathfrak{V}$ is a strict vessel that satisfies the first condition of (3.2), namely $\sigma_j \sigma_1^{-1} \sigma_k = \sigma_k \sigma_1^{-1} \sigma_j$. Then $\mathfrak{V}$ satisfies the VR conditions.*

Proof. Since the vessel is strict Φ is surjective and Φ^* is injective. We can assume without loss of generality that $\Phi \Phi^* = I_{\mathcal{E}}$. Hence from the vessel conditions we get that:

$$\begin{aligned} \sigma_j \sigma_1^{-1} \gamma_{1k} \Phi - \sigma_k \sigma_1^{-1} \gamma_{1j} \Phi - \gamma_{jk} \Phi &= \\ \sigma_j \sigma_1^{-1} (\sigma_1 \Phi A_k^* - \sigma_k \Phi A_1^*) - \sigma_k \sigma_1^{-1} (\sigma_1 \Phi A_j^* - \sigma_j \Phi A_1^*) - \sigma_j \Phi A_k^* + \sigma_k \Phi A_j^* &= \\ (\sigma_j \sigma_1^{-1} \sigma_k - \sigma_k \sigma_1^{-1} \sigma_j) \Phi A_1^*. \end{aligned}$$

Now postmultiplying by Φ^* we obtain:

$$\sigma_j \sigma_1^{-1} \gamma_{1k} - \sigma_k \sigma_1^{-1} \gamma_{1j} - \gamma_{jk} = (\sigma_j \sigma_1^{-1} \sigma_k - \sigma_k \sigma_1^{-1} \sigma_j) \Phi A_1^* \Phi^*.$$

In particular (3.3) follows from the first condition of (3.2). Next we compute:

$$\begin{aligned} \Phi^* \gamma_{1j} \sigma_1^{-1} \gamma_{1k} \Phi &= (A_j \Phi^* \sigma_1 - A_1 \Phi^* \sigma_j) \sigma_1^{-1} (\sigma_1 \Phi A_k^* - \sigma_k \Phi A_1^*) = \\ &= A_j \Phi^* \sigma_1 \Phi A_k^* - A_j \Phi^* \sigma_k \Phi A_1^* - A_1 \Phi^* \sigma_j \Phi A_k^* + A_1 \Phi^* \sigma_j \sigma_1^{-1} \sigma_k \Phi A_1^* = \\ &= \frac{1}{i} (A_j A_1 A_k^* - A_j A_1^* A_k^*) - \frac{1}{i} (A_j A_k A_1^* - A_j A_k^* A_1^*) - \frac{1}{i} (A_1 A_j A_k^* - A_1 A_j^* A_k^*) + \\ &= A_1 \Phi^* \sigma_j \sigma_1^{-1} \sigma_k \Phi A_1^* = \frac{1}{i} (A_1 A_j^* A_k^* - A_j A_k A_1^*) + A_1 \Phi^* \sigma_j \sigma_1^{-1} \sigma_k \Phi A_1^*. \end{aligned}$$

Similarly we get:

$$\Phi^* \gamma_{1k} \sigma_1^{-1} \gamma_{1j} \Phi = \frac{1}{i} (A_1 A_k^* A_j^* - A_k A_j A_1^*) + A_1 \Phi^* \sigma_k \sigma_1^{-1} \sigma_j \Phi A_1^*$$

Now since the A_j commute we obtain after premultiplying by Φ and postmultiplying by Φ^* that:

$$\gamma_{1j} \sigma_1^{-1} \gamma_{1k} - \gamma_{1k} \sigma_1^{-1} \gamma_{1j} = \Phi A_1 \Phi^* (\sigma_j \sigma_1^{-1} \sigma_k - \sigma_k \sigma_1^{-1} \sigma_j) \Phi A_1^* \Phi^*.$$

In particular the second condition of (3.2) follows from the first. Now using Proposition 4.5 we get the result. $\square$

Corollary 4.8. *Assume that $A_1, \dots, A_d$ are doubly commuting (i.e., $[A_j, A_k^*] = 0$ for every $j \neq k$) and that there exists $\xi \in \mathbb{R}^d$, such that $\sum_{j=1}^d \xi_j (A_j - A_j^*)$ is invertible when restricted to the non-Hermitian subspace, then the strict vessel they embed into satisfies the VR conditions.*

Proof. From the formulae (2.10) it follows that the σ_j commute and our assumption implies that there exists $\xi \in \mathbb{R}^d$, such that $\xi \sigma$ is invertible. Now apply Proposition 4.7 to get the result. $\square$

Remark 4.9. Let $A_1, \dots, A_d$ be a d -tuple of commuting operators, then Proposition 4.7 and Corollary 4.8 imply that the assumption that the strict vessel satisfies the VR conditions is a generalization of the doubly-commuting property.

5. SOLUTION IN THE HYPERBOLIC CASE

In this section we study the hyperbolic case and thus from now on we assume that there exists an $\epsilon > 0$, such that $\sigma_1 > \epsilon I$. Let us write again $\alpha_j = \sigma_1^{-1} \sigma_j$ and $\beta_j = \sigma_1^{-1} \gamma_{1j}$ and set $\alpha_1 = I_{\mathcal{E}}$ and $\beta_1 = 0$. In this case α_j and β_j are selfadjoint with respect to the σ_1 -inner product on $\mathcal{E}$. Without loss of generality we may assume that $\sigma_1 = I_{\mathcal{E}}$, since otherwise we can simply replace the inner product on $\mathcal{E}$ by the σ_1 -inner product and then the σ_j will be replaced by α_j and γ_{1j} will be replaced by β_j . For $x \in \mathbb{R}^d$ let us write $\alpha(x) = \sum_{j=1}^d x_j \alpha_j$ and similarly $\beta(x) = \sum_{j=1}^d x_j \beta_j$. Since $\alpha_1 = I$ there exists $\epsilon_j > 0$ small enough such that $\alpha_1 + \epsilon_j \alpha_j > \delta I$ for every $j = 2, \dots, d$ and some $\delta > 0$. Hence by changing coordinates we may assume that $\alpha_j > \delta I$ for every $j = 2, \dots, d$. The following definition describes the future cone of our system:

Definition 5.1. Let $\mathfrak{V}$ be a vessel, define the following set in $\mathbb{R}^d$:

$$\text{Pos}(\mathfrak{V}) = \{\xi \in \mathbb{R}^d \mid \exists \epsilon > 0 : \xi \sigma = \sum_{j=1}^d \xi_j \sigma_j > \epsilon I_{\mathcal{E}}\}.$$

Note that $\text{Pos}(\mathfrak{V})$ is either empty or an open convex cone in $\mathbb{R}^d$.

Recall that by a theorem of Grothendieck a function with values in $\mathcal{E}$ is smooth if and only if it is weakly smooth (cf. [19, Sec. 3.8] or [20]). Denote by $\mathcal{S}(\mathbb{R}, \mathcal{E})$ the Schwarz space of $\mathcal{E}$ -valued rapidly decreasing smooth functions on $\mathbb{R}$. Namely, $\mathcal{S}(\mathbb{R}, \mathcal{E})$ is the space of smooth $\mathcal{E}$ -valued functions, such that for every two polynomials P and Q we have that $\|P(t)Q(\frac{\partial}{\partial t})f\|$ is bounded on $\mathbb{R}$. By [42, Thm. 44.1] and [42, Ex. 44.6] we have that $\mathcal{S}(\mathbb{R}, \mathcal{E}) \cong \mathcal{S}(\mathbb{R}) \hat{\otimes} \mathcal{E}$, where the choice of the completed tensor product does not matter since $\mathcal{S}(\mathbb{R})$ is a nuclear Frechet space

(cf. [42, Ch. 51]). We also consider the space of tempered $\mathcal{E}$ -distributions on $\mathbb{R}$, namely the topological dual of $\mathcal{S}(\mathbb{R}, \mathcal{E})$, that we will denote by $\mathcal{S}'(\mathbb{R}, \mathcal{E})$. Since our goal is to discuss operators on Hilbert spaces we will use an anti-linear pairing between tempered distributions and Schwarz functions. We note that we can endow $\mathcal{S}'(\mathbb{R}, \mathcal{E})$ with the strong topology of uniform convergence on bounded subsets and that by [42, Prop. 50.7] we have $\mathcal{S}'(\mathbb{R}, \mathcal{E}) \cong \mathcal{S}'(\mathbb{R}) \hat{\otimes} \mathcal{E}$ (the identification is again anti-linear). We can also endow $\mathcal{S}'(\mathbb{R}, \mathcal{E})$ with the weak topology of pointwise convergence and those topologies will coincide if and only if $\dim \mathcal{E} < \infty$. Similarly we define $\mathcal{S}(\mathbb{R}^d, \mathcal{E})$ and $\mathcal{S}'(\mathbb{R}^d, \mathcal{E})$. We will also use the space $L^2(\mathbb{R}, \mathcal{E})$ that is the space of all weakly measurable functions $f: \mathbb{R} \rightarrow \mathcal{E}$ (this is equivalent by Pettis' theorem to strongly measurable since $\mathcal{E}$ is separable), such that:

$$\|f\|_{L^2}^2 = \int_{-\infty}^{\infty} \|f(t)\|^2 dt < \infty.$$

Similarly, if α is an invertible positive-definite operator on $\mathcal{E}$ we will define the space $L^2(\mathbb{R}, \mathcal{E}, \alpha)$ as the set of all weakly measurable functions $f: \mathbb{R} \rightarrow \mathcal{E}$, such that:

$$\int_{-\infty}^{\infty} \langle \alpha f(t), f(t) \rangle dt < \infty.$$

Then in particular we have that $L^2(\mathbb{R}, \mathcal{E}) \cong L^2(\mathbb{R}) \hat{\otimes}_H \mathcal{E}$, where $\hat{\otimes}_H$ is the tensor product of Hilbert spaces. We have a continuous embedding $\mathcal{S}(\mathbb{R}, \mathcal{E}) \hookrightarrow L^2(\mathbb{R}, \mathcal{E})$ and its image is dense. We can define the Fourier transform by considering the continuous linear map $\mathcal{F} \otimes I_{\mathcal{E}}$ on $\mathcal{S}(\mathbb{R}) \hat{\otimes} \mathcal{E}$. This is equivalent to:

$$\mathcal{F}(f)(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ist} f(s) ds.$$

Here the integral is considered as a Gel'fand-Pettis integral and by the same consideration as in the classical Plancherel theorem it extends to an isometric automorphism of $L^2(\mathbb{R}, \mathcal{E})$.

Let us assume now that the input signal is a Schwarz $\mathcal{E}$ -valued function on $\mathbb{R}^d$ and assume that the initial condition is $u = f$ on the t_1 -axis. Let us then apply the Fourier transform along the t_1 -axis to u and write $\mathcal{F}_1(u) = \hat{u}$. Then we get a system of equations ($j = 2, \dots, d$):

$$\frac{\partial \hat{u}}{\partial t_j} = i\alpha_j \tau_1 \hat{u} + i\beta_j \hat{u}.$$

Here τ_1 is the variable in the frequency domain. The initial condition is $\hat{u} = \hat{f}$ on the τ_1 -axis. Each of these equations has a solution of the form:

$$\varphi_j(\tau_1, t_2, \dots, t_d) = e^{it_j(\alpha_j \tau_1 + \beta_j)} C_j(\tau_1, t_2, \dots, t_{j-1}, t_{j+1}, \dots, t_d),$$

where C_j is an $\mathcal{E}$ -valued function. One then can proceed plugging one solution into the other equations and then using the initial condition. Since the equations are compatible and the pencils in the exponent commute by (3.2), we will get a solution:

$$\hat{u} = e^{\sum_{j=2}^d it_j(\alpha_j \tau_1 + \beta_j)} \hat{f}(\tau_1).$$

Hence a solution to the system of input compatibility equations is:

$$(5.1) \quad u(t_1, \dots, t_d) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\sum_{j=1}^d it_j(\alpha_j \tau_1 + \beta_j)} \hat{f}(\tau_1) d\tau_1.$$

This computation makes sense for Schwarz functions. We will show below (see Corollary 5.5) that for every $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ the above formula defines a smooth $\mathcal{E}$ -valued function on $\mathbb{R}^d$ (not necessarily Schwarz) that is a solution of our system. Next we would like to extend it to a wider class of functions on $\mathbb{R}$.

Note that $\mathbb{R}^d$ acts on $\mathcal{S}(\mathbb{R}, \mathcal{E})$ by $(te_j \cdot \varphi)(s) = e^{it(s\alpha_j + \beta_j)}\varphi(s)$. This function is clearly Schwarz since both α_j and β_j are selfadjoint and hence the exponent is a unitary operator on $\mathcal{E}$. We can conjugate this action by the Fourier transform, namely we get a representation of $\mathbb{R}^d$:

$$(5.2) \quad \pi(te_j)\varphi = \mathcal{F}^{-1}(e^{it(s\alpha_j + \beta_j)}\mathcal{F}(\varphi)).$$

Note that this representation is smooth by virtue of a theorem of Bruhat ([44, Prop. 4.4.1.7]) and the following fact:

$$\pi\left(\frac{\partial}{\partial t_1}\right)\varphi = \varphi'.$$

$$\pi\left(\frac{\partial}{\partial t_j}\right)\varphi = \alpha_j\varphi' + i\beta_j\varphi.$$

By [44, Prop. 4.4.1.9] we have that the contragredient representation of $\mathbb{R}^d$ on $\mathcal{S}'(\mathbb{R}, \mathcal{E})$ is also smooth, since $\mathcal{S}'(\mathbb{R}, \mathcal{E})$ is complete. Hence we get for every $f \in \mathcal{S}'(\mathbb{R}, \mathcal{E})$ a smooth function $L_f: \mathbb{R}^{d-1} \rightarrow \mathcal{S}'(\mathbb{R}, \mathcal{E})$ that is evaluated as:

$$\langle L_f(t_2, \dots, t_d)(\cdot), \varphi \rangle = \langle f, \mathcal{F}(e^{-i\sum_{j=2}^d t_j(s\alpha_j + \beta_j)}\mathcal{F}^{-1}(\varphi)) \rangle.$$

Note that if $u \in \mathcal{S}(\mathbb{R}^d, \mathcal{E})$, that solves the system of input compatibility conditions and f is its restriction to the t_1 -axis, then (5.1) implies that $u(\cdot, t_2, \dots, t_d) = L_f(t_2, \dots, t_d)(\cdot)$. Note also, that in general the function L_f solves the input compatibility conditions in the following sense:

$$\pi\left(\frac{\partial}{\partial t_j}\right)L_f = \alpha_j L_{f'} + i\beta_j L_f = (\alpha_j \pi\left(\frac{\partial}{\partial t_1}\right) + i\beta_j)L_f.$$

Remark 5.2. Recall that equation (3.3) implies that the remaining equations are satisfied, if those involving σ_1 are satisfied.

Proposition 5.3. *The function L_f defines a tempered $\mathcal{E}$ -valued distribution u_f on $\mathbb{R}^d$ as follows:*

$$(5.3) \quad \langle u_f, \psi \rangle = \int_{\mathbb{R}^{d-1}} \langle L_f(t_2, \dots, t_d)(\cdot), \psi(\cdot, t_2, \dots, t_d) \rangle dt_2 \cdots dt_d.$$

Here $\psi \in \mathcal{S}(\mathbb{R}^d, \mathcal{E})$ is a Schwarz function.

Proof. We note that since $f \in \mathcal{S}'(\mathbb{R}, \mathcal{E})$ there exist polynomials $P(t)$ and $Q(t)$ and a constant $C > 0$, such that for every choice of $t_2, \dots, t_d$ and ψ , we have:

$$\begin{aligned}
|\langle L_f(t_2, \dots, t_d)(\cdot), \psi(\cdot, t_2, \dots, t_d) \rangle| &= \\
|\langle f, \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\psi(\cdot, t_2, \dots, t_d))) \rangle| &\leq \\
C \sup_{t_1 \in \mathbb{R}} \|P(t_1) Q(\frac{\partial}{\partial t_1}) \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\psi(t_1, t_2, \dots, t_d)))\| &= \\
C \sup_{t_1 \in \mathbb{R}} \|\mathcal{F}(P(i \frac{\partial}{\partial s}) Q(-is) e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\psi(t_1, t_2, \dots, t_d)))\| &= \\
C \sup_{t_1 \in \mathbb{R}} \|\mathcal{F}(P(i \frac{\partial}{\partial s}) e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(Q(\frac{\partial}{\partial t_1}) \psi(t_1, t_2, \dots, t_d)))\| &\leq \\
CC' \sup_{s \in \mathbb{R}} \|P(i \frac{\partial}{\partial s}) e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}((1 + \frac{\partial}{\partial t_1^2}) Q(\frac{\partial}{\partial t_1}) \psi(t_1, t_2, \dots, t_d))\|. &
\end{aligned}$$

The last inequality is due to the fact that for every function $\varphi \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ we have that:

$$(5.4) \quad \sup_{s \in \mathbb{R}} \|\mathcal{F}(\varphi)(s)\| \leq C' \sup_{s \in \mathbb{R}} \|(1 + s^2)\varphi(s)\|.$$

For the derivative of the exponent applied to a Schwarz function we have the following bound (here we write $\eta = \mathcal{F}^{-1}((1 + \frac{\partial}{\partial t_1^2}) Q(\frac{\partial}{\partial t_1}) \psi(t_1, t_2, \dots, t_d))$):

$$\left\| \frac{\partial}{\partial s} \left(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \eta(s) \right) \right\| \leq \left\| \frac{\partial}{\partial s} \left(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \right) \eta(s) \right\| + \left\| \frac{\partial \eta}{\partial s} \right\|$$

We use here the fact that the exponent is a unitary operator for every choice of real s . Now we have the following well known equality:

$$\frac{\partial}{\partial s} \left(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \right) = - \int_0^1 e^{-i w \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \left(\sum_{j=2}^d t_j \alpha_j \right) e^{-(1-w) i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} dw.$$

Thus we obtain the inequality:

$$\left\| \frac{\partial}{\partial s} \left(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \eta(s) \right) \right\| \leq \sum_{j=2}^d |t_j| \|\alpha_j\| \|\eta(s)\| + \left\| \frac{\partial \eta}{\partial s} \right\|.$$

Notice that in the expression $\frac{\partial \eta}{\partial s}$ we can push $\frac{\partial}{\partial s}$ into $\mathcal{F}^{-1}$ replacing it by $-it_1$. Using estimate (5.4) we can then get rid of the Fourier transform. Applying these considerations to every monomial in P we eventually get that there exist a constant $\tilde{C} > 0$ and polynomials $\tilde{P}, \tilde{Q} \in \mathbb{C}[t_1, \dots, t_d]$, such that:

$$|\langle L_f(t_2, \dots, t_d)(\cdot), \psi(\cdot, t_2, \dots, t_d) \rangle| \leq \tilde{C} \sup_{t_1 \in \mathbb{R}} \|\tilde{P}(t) \tilde{Q}(\frac{\partial}{\partial t}) \psi(t)\|.$$

As for the integral we write:

$$\begin{aligned}
\left| \int_{\mathbb{R}^{d-1}} \langle L_f(t_2, \dots, t_d)(\cdot), \psi(\cdot, t_2, \dots, t_d) \rangle dt_2 \cdots dt_d \right| &\leq \\
C_0 \sup_{t=(t_1, \dots, t_d) \in \mathbb{R}^d} \|(1 + t_2^2 + \dots + t_d^2)^d \tilde{P}(t) \tilde{Q}(\frac{\partial}{\partial t}) \psi(t)\|. &
\end{aligned}$$

Here $C_0 = \tilde{C} \int_{\mathbb{R}^{d-1}} (1 + t_2^2 + \dots + t_d^2)^{-d} dt_2 \dots t_d$. $\square$

Proposition 5.4. *The tempered distribution u_f defined in (5.3) is a weak solution for the system of input compatibility conditions.*

Proof. We want to show that for every $f \in \mathcal{S}'(\mathbb{R}, \mathcal{E})$ and every $\psi \in \mathcal{S}(\mathbb{R}, \mathcal{E})$, we have:

$$\left\langle \frac{\partial u_f}{\partial t_k}, \psi \right\rangle = \left\langle \alpha_k \frac{\partial u_f}{\partial t_1} + i\beta_k u_f, \psi \right\rangle.$$

Since α_k and β_k are selfadjoint, by taking the adjoint we see that the desired equality becomes:

$$\left\langle u_f, \frac{\partial \psi}{\partial t_k} \right\rangle = \left\langle u, \alpha_k \frac{\partial \psi}{\partial t_1} + i\beta_k \psi \right\rangle.$$

For $\psi \in \mathcal{S}(\mathbb{R}^d, \mathcal{E})$, we want to compute $\left\langle u_f, \frac{\partial}{\partial t_k} \psi \right\rangle$, for some $k = 2, \dots, d$. For fixed $t_2, \dots, t_d \in \mathbb{R}^d$ we have:

$$\left\langle L_f(t_2, \dots, t_d)(\cdot), \frac{\partial}{\partial t_k} \psi \right\rangle = \left\langle f, \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\frac{\partial}{\partial t_k} \psi)) \right\rangle.$$

Now we compute:

$$\begin{aligned} \frac{\partial}{\partial t_k} \left(\mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\psi)) \right) &= \\ \mathcal{F} \left(\frac{\partial}{\partial t_k} \left(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \right) \mathcal{F}^{-1}(\psi) \right) &+ \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\frac{\partial \psi}{\partial t_k})) = \\ - \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} (is\alpha_k + i\beta_k) \mathcal{F}^{-1}(\psi)) &+ \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\frac{\partial \psi}{\partial t_k})) = \\ - \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\alpha_k \frac{\partial \psi}{\partial t_1} + i\beta_k \psi)) &+ \mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\frac{\partial \psi}{\partial t_k})). \end{aligned}$$

Now we apply f and integrate on $\mathbb{R}^{d-1}$ to get:

$$\begin{aligned} \int_{\mathbb{R}^{d-1}} \left\langle f, \frac{\partial}{\partial t_k} \left(\mathcal{F}(e^{-i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \mathcal{F}^{-1}(\psi(\cdot, t_2, \dots, t_d))) \right) \right\rangle &= \\ - \left\langle f u_f, \alpha_k \frac{\partial \psi}{\partial t_1} + i\beta_k \psi \right\rangle &+ \left\langle u_f, \frac{\partial \psi}{\partial t_k} \right\rangle. \end{aligned}$$

It remains to note that the left hand side is zero since ψ is a Schwarz function on $\mathbb{R}^d$ and f is a distribution on $\mathbb{R}$. $\square$

Lemma 5.5. *If $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ then u_f is a smooth function on $\mathbb{R}^d$ that solves the system of input compatibility equations. Furthermore, u_f is given by the formula (5.1).*

Proof. Let $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ and note that by definition:

$$L_f(t_2, \dots, t_d)(\cdot) = (\pi(\cdot, t_2, \dots, t_d)f)(0).$$

Here π is the representation defined in (5.2). Therefore, the associated u is the smooth function given by (5.1) and since it is a weak solution it is a solution. $\square$

Lemma 5.6. *The representation π (defined by (5.2)) of $\mathbb{R}^d$ on $\mathcal{S}(\mathbb{R}, \mathcal{E})$ extends to a unitary representation of $\mathbb{R}^d$ on $L^2(\mathbb{R}, \mathcal{E})$.*

Proof. Since both α_j and β_j are selfadjoint for every $j = 1, \dots, d$ and t_j and s are real, the multiplication by $e^{i \sum_{j=1}^d t_j (s\alpha_j + \beta_j)}$ is a unitary operator. $\square$

To better understand the solutions we will study their behavior on lines. Consider the formula (5.1). If we fix a line $\tau x + y$ in $\mathbb{R}^d$, then we define for every $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ the following operator:

$$(5.5) \quad (\Lambda(x, y)f)(\tau) = (\pi(\tau x + y)f)(0) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i\tau(s\alpha(x) + \beta(x))} e^{i(s\alpha(y) + \beta(y))} \hat{f}(s) ds.$$

Notice that by Lemma 5.5 we have that u_f is defined by the formula (5.1) and thus, for every $x, y \in \mathbb{R}^d$ and $\tau \in \mathbb{R}$, we have:

$$(5.6) \quad u_f(\tau x + y) = (\Lambda(x, y)f)(\tau).$$

Let us summarize the discussion above in the following theorem:

Theorem 5.7. *Given a vessel $\mathfrak{V}$ satisfying the VR conditions and such that $\sigma_1 > \epsilon I$ for some $\epsilon > 0$, for every initial condition $f \in \mathcal{S}'(\mathbb{R}, \mathcal{E})$ we have a weak solution of the system of input compatibility conditions in tempered distributions defined by (5.3). Furthermore, the following holds:*

- If $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$, then u_f is a smooth function on $\mathbb{R}^d$, that solves the system of input compatibility equations.
- Fix $x \in \mathbb{R}^d$, such that $\alpha(x) > \epsilon I$ as well, then for every $y \in \mathbb{R}^d$, we can extend $\Lambda(x, y)$ to an isometric isomorphism from $L^2(\mathbb{R}, \mathcal{E})$ to $L^2(\mathbb{R}, \mathcal{E}, \alpha(x))$. Furthermore, for $x, x', y \in \mathbb{R}^d$, the map $\Lambda(x, y)\Lambda(x', y)^*$ is a causal isometric isomorphism, i.e., for every $f \in L^2(\mathbb{R}, \mathcal{E})$ we have the following equalities:

$$(5.7) \quad \begin{aligned} \int_0^\infty \langle \alpha(x)(\Lambda(x, y)f)(s), (\Lambda(x, y)f)(s) \rangle ds &= \int_0^\infty \langle \alpha(x')(\Lambda(x', y)f)(s), (\Lambda(x', y)f)(s) \rangle ds, \\ \int_{-\infty}^0 \langle \alpha(x)(\Lambda(x, y)f)(s), (\Lambda(x, y)f)(s) \rangle ds &= \int_{-\infty}^0 \langle \alpha(x')(\Lambda(x', y)f)(s), (\Lambda(x', y)f)(s) \rangle ds. \end{aligned}$$

In particular if $y = 0$ and $x' = e_1$, then we get that $\Lambda(x, 0)$ is a causal isometric isomorphism from $L^2(\mathbb{R}, \mathcal{E})$ to $L^2(\mathbb{R}, \mathcal{E}, \alpha(x))$.

- If f is a twice continuously differentiable function on $\mathbb{R}$, such that $f, f', f'' \in L^2(\mathbb{R}, \mathcal{E})$, then u_f is a locally integrable function given by the formula (5.1) and for every $x, y \in \mathbb{R}^d$, such that $x \in \text{Pos}(\mathfrak{V})$, the restriction of u_f to the line $\tau x + y$ is given by $\Lambda(x, y)f$, namely:

$$u_f(\tau x + y) = (\Lambda(x, y)f)(\tau).$$

- If f is a twice continuously differentiable function on $\mathbb{R}$, such that $f, f', f'' \in L^2(\mathbb{R}, \mathcal{E})$, then u_f is a C^1 , $\mathcal{E}$ -valued function on $\mathbb{R}^d$ that solves the input compatibility conditions. Furthermore, u_f is uniquely determined by its restriction to the t_1 -axis (or in fact any line with direction vector in $\text{Pos}(\mathfrak{V})$).
- If $f \in L^2(\mathbb{R}, \mathcal{E})$ then for every $\xi \in \text{Pos}(\mathfrak{V})$ and every $\psi \in \mathcal{S}(\mathbb{R}^d, \mathcal{E})$ we have that:

$$\langle u_f, \psi \rangle = \int_{\xi^\perp} \int_{-\infty}^{\infty} \langle (\Lambda(\xi, \eta)f)(s), \psi(\xi s + \eta) \rangle ds d\eta.$$

Proof. We have already proved the first claim of the theorem (see Lemma 5.5). To prove the second we note that by virtue of Lemma 5.5, if $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ then $\Lambda(x, y)f$ is a smooth function on $\mathbb{R}$ and is a restriction of a solution for the system of input compatibility equations to the line $\tau x + y$ (see (5.6)). Now we recall that $\alpha(x) > 0$ and hence we can apply [11, Prop. 2.1] ((the proof given there in the case $d = 2$ extends verbatim to the case of an arbitrary d) to get that the equations (5.7) hold in this case. Now we can extend $\Lambda(x, y)$ as an isometry from $L^2(\mathbb{R}, \mathcal{E})$ to $L^2(\mathbb{R}, \mathcal{E}, \alpha(x))$.

We will prove the third and the fourth claims together. The proof follows similar lines to [14, Thm. 7.3.5]. We note that by assumption there exists a function $g \in L^2(\mathbb{R}, \mathcal{E})$ and a constant $C > 0$, such that for every $s \in \mathbb{R}$, we have:

$$\|\hat{f}(s)\| \leq C(1 + |s|^2)^{-1} \|\hat{g}(s)\|.$$

Hence for every $t \in \mathbb{R}^d$, we have:

$$\|e^{i \sum_{j=1}^d t_j (s\alpha_j + \beta_j)} \hat{f}(s)\| \leq C(1 + |s|^2)^{-1} \|\hat{g}(s)\|.$$

We conclude that:

$$\begin{aligned} \int_{-\infty}^{\infty} \|e^{i \sum_{j=1}^d t_j (s\alpha_j + \beta_j)} \hat{f}(s)\| ds &\leq C \int_{-\infty}^{\infty} (1 + |s|^2)^{-1} \|\hat{g}(s)\| ds \leq \\ &C \sqrt{\int_{-\infty}^{\infty} \|\hat{g}(s)\|^2 ds} \sqrt{\int_{-\infty}^{\infty} (1 + |s|^2)^{-2} ds} < \infty. \end{aligned}$$

Hence we have $L_f(t_2, \dots, t_d)(t_1) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i \sum_{j=1}^d t_j (s\alpha_j + \beta_j)} \hat{f}(s) ds$. Thus u_f is just integrating L_f against φ , and therefore we can conclude that u_f is a function and we can identify $u_f(t_1, \dots, t_d) = L_f(t_2, \dots, t_d)(t_1)$. Furthermore, it is now clear that the restriction of u_f to lines is given by $\Lambda(x, y)f$.

Let now $0 < |h| < 1$, we write:

$$\begin{aligned} \frac{u_f(t_1 + h, t_2, \dots, t_d) - u_f(t_1, \dots, t_d)}{h} &= \\ &\frac{1}{\sqrt{2\pi}h} \int_{-\infty}^{\infty} e^{i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} \left(e^{is(t_1+h)} - e^{ist_1} \right) \hat{f}(s) ds \end{aligned}$$

Since $e^{is(t_1+h)} - e^{ist_1} = is \int_{t_1}^{t_1+h} e^{isx} dx$, we can use the integral mean value theorem to obtain:

$$e^{is(t_1+h)} - e^{ist_1} = ish e^{isc}.$$

Here c lies between t_1 and $t_1 + h$. Therefore:

$$\frac{u_f(t_1 + h, t_2, \dots, t_d) - u_f(t_1, \dots, t_d)}{h} = \frac{i}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i \sum_{j=2}^d t_j (s\alpha_j + \beta_j)} s e^{isc} \hat{f}(s) ds$$

The integral thus converges, since $\|s\hat{f}(s)\| \leq C|s|(1 + |s|^2)^{-1} \|\hat{g}(s)\|$ and the same argument as above applies. Now applying the dominated convergence theorem we can deduce that $\frac{\partial u_f}{\partial t_1}$ exists and is continuous. A similar argument applies to every t_j .

Uniqueness follows from the fact that we can consider the lines parallel to the t_1 -axis and use the second part to note that if a solution is 0 on the t_1 -axis then using the isometry it is 0 along every such line and thus it is identically zero.

If f is a Schwartz function, then u_f is a smooth function and we have the desired equality by the definition of u_f and Fubini's theorem. Now we approximate $f \in L^2(\mathbb{R}, \mathcal{E})$ be a sequence of Schwartz functions and since $\Lambda(\xi, \eta)$ is an isometry from $L^2(\mathbb{R}, \mathcal{E})$ to $L^2(\mathbb{R}, \mathcal{E}, \alpha(\xi))$, we get that $\|\Lambda(\xi, \eta)f\| \leq C(\xi)\|f\|$. Thus for M sufficiently large:

$$\begin{aligned} & \left\| \int_{\xi^\perp} \int_{-\infty}^{\infty} \langle (\Lambda(\xi, \eta)(f - f_n))(s), \psi(\xi s + \eta) \rangle ds d\eta \right\| \leq C(\xi) \int_{\xi^\perp} \|f - f_n\| \|\psi(\xi s + \eta)\|_{L^2} d\eta \\ & = C(\xi) \int_{\xi^\perp} \|f - f_n\| (1 + \|\eta\|^{2M})^{-1} \|(1 + \|\eta\|^{2M})\psi(\xi s + \eta)\|_{L^2} d\eta \leq C'(\xi) \|f - f_n\|. \end{aligned}$$

The last inequality follows from the fact that:

$$\begin{aligned} \|(1 + \|\eta\|^{2M})\psi(\xi s + \eta)\|_{L^2}^2 &= \int_{-\infty}^{\infty} (1 + s^2)^{-1} \|(1 + \|\eta\|^{2M})(1 + s^2)\psi(\xi s + \eta)\|_{L^2}^2 ds \\ &\leq \sup_{(s, \eta) \in \mathbb{R}^d} \|(1 + \|\eta\|^{2M})(1 + s^2)\psi(\xi s + \eta)\|_{L^2}^2 \int_{-\infty}^{\infty} (1 + s^2)^{-1} ds. \end{aligned}$$

Now applying the same consideration to the case when $\xi = e_1$ we get that for every $\psi \in \mathcal{S}(\mathbb{R}^d, \mathcal{E})$ we have $\langle u_{f_n}, \psi \rangle \rightarrow \langle u_f, \psi \rangle$ and additionally:

$$\begin{aligned} \langle u_f, \psi \rangle &= \lim_{n \rightarrow \infty} \langle u_{f_n}, \psi \rangle = \lim_{n \rightarrow \infty} \int_{\xi^\perp} \int_{-\infty}^{\infty} \langle (\Lambda(\xi, \eta)f_n)(s), \psi(\xi s + \eta) \rangle ds d\eta \\ &= \int_{\xi^\perp} \int_{-\infty}^{\infty} \langle (\Lambda(\xi, \eta)f)(s), \psi(\xi s + \eta) \rangle ds d\eta \end{aligned}$$

□

Remark 5.8. In fact, the third and fourth statements of the preceding theorem is true for functions in the Sobolev space $W^{2,2}(\mathbb{R}, \mathcal{E})$. For more details on Sobolev spaces of Banach space valued functions, see [2].

Remark 5.9. Note that for every $f \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ it is immediate from (5.5) and the definition of π that for every $t \in \mathbb{R}^d$:

$$\Lambda(x, y)(\pi(t)f) = \Lambda(x, y + t)(f)$$

Thus, in particular, if $x = e_j$ one of the vectors in the standard basis of $\mathbb{R}^d$, then:

$$(\Lambda(e_j, 0)(\pi(t_j e_j)f))(\tau) = (\Lambda(e_j, t_j e_j)f)(\tau) = (\Lambda(e_j, 0)f)(\tau + t_j).$$

The last equality follows from Equation (5.5). We conclude that if $\alpha_j > \epsilon I$ for some $\epsilon > 0$, then $\Lambda_j = \Lambda(e_j, 0)$ intertwines the action of $\mathbb{R}$ on $L^2(\mathbb{R}, \mathcal{E}, \alpha_j)$ by translations with the action of $\mathbb{R}$ on $L^2(\mathbb{R}, \mathcal{E})$ by the restriction of π to the one parameter subgroup generated by e_j .

6. UNITARY DILATION OF SEMIGROUPS

In order to apply the results of the previous section to dilation theory we need a definition

Definition 6.1. Let $\mathcal{A} = (A_1, \dots, A_d)$ be a d -tuple of commuting dissipative operators on a separable Hilbert space $\mathcal{H}$. We say that $\mathcal{A}$ has the dissipative embedding property if $\mathcal{A}$ can be embedded in a vessel $\mathfrak{V}$ satisfying the VR conditions and such that $\text{Pos}(\mathfrak{V}) \neq \emptyset$ and for every $j = 1, \dots, d$ the standard basis vector $e_j \in \overline{\text{Pos}(\mathfrak{V})}$.

Lemma 6.2. Assume that $\mathcal{A}$ is a d -tuple of commuting dissipative operators on a separable Hilbert space $\mathcal{H}$ and assume that we can embed $\mathcal{A}$ in a vessel $\mathfrak{V}$, such that for every $j = 1, \dots, d$, we have $\sigma_j \geq 0$. Then $\sum_{j=1}^d \text{Im } \sigma_j = \mathcal{E}$ if and only if $\text{Pos}(\mathfrak{V})$ is not empty. In that case we have that $\mathbb{R}_{\geq 0}^d \subseteq \text{Pos}(\mathfrak{V})$. In particular if we can embed $\mathcal{A}$ in a vessel $\mathfrak{V}$, such that for every $j = 1, \dots, d$, we have $\sigma_j \geq 0$, $\sum_{j=1}^d \text{Im } \sigma_j = \mathcal{E}$ and the vessel satisfies the VR conditions, then $\mathcal{A}$ has the dissipative embedding property.

Proof. If there is a point $\xi \in \text{Pos}(\mathfrak{V})$, then there exists $\epsilon > 0$, such that for every $u \in \mathcal{E}$, we have:

$$\sum_{j=1}^n \xi_j \langle \sigma_j u, u \rangle \geq \epsilon \|u\|^2.$$

Since each σ_j is positive semi-definite, if we omit the terms with $\xi_j \leq 0$ from the above sum, then we just increase it. Hence:

$$\sum_{\xi_j > 0} \xi_j \langle \sigma_j u, u \rangle \geq \epsilon \|u\|^2.$$

This implies that the operator $\sum_{\xi_j > 0} \xi_j \sigma_j \geq \epsilon I$ and is in particular invertible, hence $\sum_{j=1}^d \text{Im } \sigma_j = \mathcal{E}$.

Now assume conversely that $\sum_{j=1}^d \text{Im } \sigma_j = \mathcal{E}$. Since the σ_j are positive semi-definite, they admit selfadjoint square roots. We have that for every $j = 1, \dots, d$, $\text{Im } \sigma_j \subseteq \text{Im } \sqrt{\sigma_j}$ and hence $\sum_{j=1}^d \text{Im } \sqrt{\sigma_j} = \mathcal{E}$. We conclude that the row $(\sqrt{\sigma_1}, \dots, \sqrt{\sigma_d})$ is strictly surjective, therefore there exists $\epsilon > 0$, such that for every $u \in \mathcal{E}$:

$$\left\| \begin{pmatrix} \sqrt{\sigma_1} u \\ \vdots \\ \sqrt{\sigma_d} u \end{pmatrix} \right\|^2 \geq \epsilon \|u\|^2.$$

But since $\|\sqrt{\sigma_j} u\|^2 = \langle \sigma_j u, u \rangle$, we have that the point $(1, \dots, 1) \in \text{Pos}(\mathfrak{V})$.

From the above discussion we have that $\text{Pos}(\mathfrak{V}) \cap \mathbb{R}_{>0}^d \neq \emptyset$. On the other hand for every $\xi \in \mathbb{R}_{>0}^d$, we replace σ_j with $\sqrt{\xi_j} \sigma_j$ in the above argument and get that the point $\xi \in \text{Pos}(\mathfrak{V})$. $\square$

Theorem 6.3. If a d -tuple $\mathcal{A}$ of commuting dissipative operators on a separable Hilbert space $\mathcal{H}$ has the dissipative embedding property, then the semigroup of contractions generated by $\mathcal{A}$ admits a commutative unitary dilation.

This theorem is a corollary of the following slightly more general theorem, that we will prove in the next section.

Theorem 6.4. Assume that $\mathcal{A}$ is a d -tuple of commuting dissipative operators on a Hilbert space and assume that $\mathcal{A}$ can be embedded into a vessel $\mathfrak{V}$ that satisfies the VR conditions and $\text{Pos}(\mathfrak{V}) \neq \emptyset$, then the semigroup of contractions generated by $\mathcal{A}$ restricted to $\overline{\text{Pos}(\mathfrak{V})}$ admits a commutative unitary dilation.

Proof of Theorem 6.3. We have that $\mathbb{R}_{\geq 0}^d \subset \overline{\text{Pos}(\mathfrak{V})}$ (in fact by Lemma 6.2 $\mathbb{R}_{>0}^d \subset \text{Pos}(\mathfrak{V})$) and thus our semigroup admits a unitary dilation by Theorem 6.4. $\square$

We shall now deduce a few corollaries from Theorem 6.3. We first note that we have the following weak form of Ando's theorem (see also [32], [38] and [39]):

Corollary 6.5. *Let A_1 and A_2 be two commuting dissipative operators, such that $\text{Im}(A_1 - A_1^*) + \text{Im}(A_2 - A_2^*)$ is closed. Then the semigroup they generate admits a commutative unitary dilation*

Proof. Note that every vessel of a pair of commuting operators satisfies the VR conditions vacuously. Furthermore, by our assumption we can embed A_1 and A_2 into a strict vessel that satisfies the conditions of Lemma 6.2. Therefore, they have the dissipative embedding property and we are done. $\square$

Recall that given a strongly continuous one-parameter semigroup C of contractions on a Hilbert space $\mathcal{H}$, by a theorem of Hille and Yosida, it has a generator, namely $C(t) = e^{iAt}$, where A is a dissipative (generally unbounded) operator on $\mathcal{H}$. If we apply the Cayley transform to A we obtain a contractive operator $T = (A - iI)(A + iI)^{-1}$, that is called the cogenerator of the semigroup. Note that the semigroup can be recovered from T via exponentiation of the inverse Cayley transform, namely $C(t) = \exp(t(T - I)(T + I)^{-1})$. We can also recover the cogenerator from the semigroup directly by the following formula (see [41] for details):

$$(6.1) \quad T = \lim_{s \rightarrow 0+} \varphi_s(C(s)),$$

$$(6.2) \quad \varphi_s(\lambda) = \frac{\lambda - 1 + s}{\lambda - 1 - s} = \frac{1 - s}{1 + s} - \frac{2s}{1 + s} \sum_{n=1}^{\infty} \frac{\lambda^n}{(1 + s)^n}.$$

Furthermore, it was proved by Sz.-Nagy that T is unitary if and only if C is a unitary semigroup. Now if we have a multi-parameter commutative group of unitaries, its generators are strongly commuting selfadjoint operators (in the sense that the associated projection valued measures commute). Therefore, applying the Cayley transform we get a commuting d -tuple of unitaries. Using (6.1) and (6.2) we conclude that for a commutative semigroup of contractions that admit a commutative unitary dilation, the cogenerators of the unitary group are commuting dilations of the cogenerators of the original semigroup. This discussion leads us to the following negative result:

Proposition 6.6. *Not every d -tuple of commuting dissipative operators on a separable Hilbert space has the dissipative embedding property.*

Proof. Consider the Parrot example described in [29]. The example is three commuting contractive matrices with spectrum concentrated at 0. Hence we can apply the Cayley transform to obtain three commuting dissipative operators, $\mathcal{A} = (A_1, A_2, A_3)$. If $\mathcal{A}$ had the dissipative embedding property then by Theorem 6.3 the semigroup they generate would have had a commutative unitary dilation. Thus the cogenerators of this dilation would have been commuting unitary dilations of the original Parrot example and that is a contradiction. $\square$

The following corollary is also well known, see for example [32] and [35].

Corollary 6.7. *If $A_1, \dots, A_d$ are doubly commuting, dissipative operators on $\mathcal{H}$, such that $\sum_{j=1}^d \text{Im}(A_j - A_j^*)$ is closed, then the semigroup they generate admits a dilation to a commutative semigroup of unitaries.*

Proof. Note that since $\sum_{j=1}^d \text{Im}(A_j - A_j^*)$ is closed, by Lemma 6.2 we have that $\text{Pos}(\mathfrak{V})$ contains the positive orthant, where $\mathfrak{V}$ is the strict vessel. By Corollary 4.8

we have that $\mathfrak{V}$ satisfies the VR conditions and thus we can apply Theorem 6.3 to get the result. $\square$

7. CONSTRUCTION OF THE DILATION (PROOF OF THEOREM 6.4)

We are given a d -tuple of dissipative operators $\mathcal{A} = (A_1, \dots, A_d)$ on a separable Hilbert space $\mathcal{H}$ embedded in a commutative vessel $\mathfrak{V}$ that satisfies the VR conditions and such that $\text{Pos}(\mathfrak{V}) \neq \emptyset$. We will construct a Hilbert space $\mathcal{K}$, an isometric embedding $\iota: \mathcal{H} \hookrightarrow \mathcal{K}$ and a unitary representation $\rho: \mathbb{R}^d \rightarrow B(\mathcal{K})$, such that for every $t \in \text{Pos}(\mathfrak{V})$ and every $h \in \mathcal{H}$ we have $\iota^* \rho(t) \iota(h) = e^{itA} h$. Notice that by passing to the SOT-limit we see that the result still holds for $t \in \overline{\text{Pos}(\mathfrak{V})}$.

We assume without loss of generality that $e_1 \in \text{Pos}(\mathfrak{V})$. As in Section 5 we may (changing the inner product on $\mathcal{E}$) assume that $\sigma_1 = I_{\mathcal{E}}$. Recall from [27] and [36, Prop. 1.3.1] that given a $u \in C^1(\mathbb{R}^d, \mathcal{E})$ that satisfies the input compatibility conditions, we can solve the time domain system of equations for any initial condition $x(0) = h \in \mathcal{H}$ using formula (2.7). In particular:

$$(7.1) \quad x(t) = x(t, 0, \dots, 0) = e^{itA_1} \left(h - i \int_0^t e^{-isA_1} \Phi^* u(s) ds \right).$$

This idea allows us to decompose the space of "nice" trajectories of the associated system into a direct sum of the form $\mathcal{W}_{out} \oplus \mathcal{H} \oplus \mathcal{W}_{in}$. Here $\mathcal{H}$ represents the initial condition. We then introduce using the theory developed in Section 5 a unitary representation of $\mathbb{R}^d$ on this space, such that the compression of its $\text{Pos}(\mathfrak{V})$ semigroup to $\mathcal{H}$ is our initial semigroup of contractions restricted to $\text{Pos}(\mathfrak{V})$.

Let us consider first the case of a single operator with $\sigma_1 = I_{\mathcal{E}}$. This is a classical construction one can find for example in [23], [24], [41], [30] and [34]. Note that in this case the one-parameter semigroup $T(t) = e^{itA_1}$ for $t > 0$ is a semigroup of contractions on $\mathcal{H}$. Set $\mathcal{K} = L^2(\mathbb{R}_{<0}, \mathcal{E}) \oplus \mathcal{H} \oplus L^2(\mathbb{R}_{>0}, \mathcal{E})$ and we are going to describe a dilation of T to $\mathcal{K}$. To do this we need the following lemma:

Lemma 7.1. *Given a triple $(y, h, u) \in \mathcal{K}$ there exists a unique (strongly) absolutely continuous function $x: \mathbb{R} \rightarrow \mathcal{H}$, such that for $t > 0$:*

$$ix'(t) + A_1 x(t) = \Phi^* u(t)$$

and for $t < 0$:

$$ix'(t) + A_1^* x(t) = \Phi^* y(t).$$

We then extend y to $\tilde{y}: \mathbb{R} \rightarrow \mathcal{E}$ by defining for $t > 0$:

$$\tilde{y}(t) = u(t) - i\Phi x(t)$$

and we extend u to $\tilde{u}: \mathbb{R} \rightarrow \mathcal{E}$ by defining for $t < 0$:

$$\tilde{u}(t) = y(t) + i\Phi x(t).$$

Then we have $\tilde{y}, \tilde{u} \in L^2(\mathbb{R}, \mathcal{E})$.

Proof. We define x in terms of u using (7.1) for $t > 0$ and in terms of y using the analog of (7.1) for the adjoint system (see (2.8)) for $t < 0$:

$$(7.2) \quad x(t) = \begin{cases} e^{itA_1} \left(h - i \int_0^t e^{-isA_1} \Phi^* u(s) ds \right), & t > 0; \\ e^{itA_1^*} \left(h + i \int_t^0 e^{-isA_1^*} \Phi^* y(s) ds \right), & t < 0; \end{cases}$$

Then clearly x is an absolutely continuous $\mathcal{H}$ -valued function on $\mathbb{R}$. Now from the energy conservation equations (2.9) we get that for $t > 0$:

$$\|x(t)\|^2 - \|h\|^2 = \int_0^t \langle \tilde{u}, \tilde{u} \rangle - \int_0^t \langle \tilde{y}, \tilde{y} \rangle.$$

We conclude that:

$$\int_0^t \langle \tilde{y}, \tilde{y} \rangle \leq \int_0^t \langle \tilde{u}, \tilde{u} \rangle + \|h\|^2.$$

Therefore $\tilde{y} \in L^2(\mathbb{R}, \mathcal{E})$ and similarly for $\tilde{u}$. $\square$

Notice that the trajectory $(\tilde{y}, x, \tilde{u})$ is the unique trajectory of the system (equivalently $(\tilde{u}, x, \tilde{y})$ is a unique trajectory of the adjoint system), such that $\tilde{y}|_{\mathbb{R}_{<0}} = y$, $x(0) = h$ and $\tilde{u}|_{\mathbb{R}_{>0}} = u$. The following proposition provides a dilation of the one parameter semigroup of contractions generated by A_1 .

Proposition 7.2. *There exists a unitary representation ρ of the Lie group $\mathbb{R}$ on $\mathcal{K}$, such that if P is the projection onto $\mathcal{H}$, then:*

$$P\rho(t)(0, h, 0) = (0, e^{itA_1}h, 0).$$

Proof. Let $(y, h, u) \in \mathcal{K}$, let $(\tilde{y}, x, \tilde{u})$ be the unique trajectory of the system associated to our triple, where x is defined by (7.2). Denote by $\tilde{y}_t(s) = \tilde{y}(s+t)$ and similarly $\tilde{u}_t(s) = \tilde{u}(s+t)$, for every $t \in \mathbb{R}$. Now we define our representation as follows:

$$\rho(t)(y, h, u) = (\tilde{y}_t|_{\mathbb{R}_{<0}}, x(t), \tilde{u}_t|_{\mathbb{R}_{>0}}).$$

Using the energy balance equations we obtain for $t > 0$:

$$\|x(t)\|^2 = \|x(0)\|^2 + \int_0^t \langle u(s), u(s) \rangle ds - \int_0^t \langle \tilde{y}(s), \tilde{y}(s) \rangle ds.$$

Therefore for $t > 0$:

$$\begin{aligned} \|\rho(t)(y, h, u)\|^2 &= \|x(t)\|^2 + \int_{-\infty}^0 \langle \tilde{y}_t(s), \tilde{y}_t(s) \rangle ds + \int_0^\infty \langle \tilde{u}_t(s), \tilde{u}_t(s) \rangle ds = \\ &= \|x(t)\|^2 + \int_{-\infty}^0 \langle \tilde{y}(s+t), \tilde{y}(s+t) \rangle ds + \int_0^\infty \langle \tilde{u}(s+t), \tilde{u}(s+t) \rangle ds = \\ &= \|h\|^2 + \int_0^t \langle u(s), u(s) \rangle ds - \int_0^t \langle \tilde{y}(s), \tilde{y}(s) \rangle ds + \int_{-\infty}^t \langle \tilde{y}(s), \tilde{y}(s) \rangle ds + \\ &+ \int_t^\infty \langle \tilde{u}(s), \tilde{u}(s) \rangle ds = \|h\|^2 + \int_0^\infty \langle u(s), u(s) \rangle ds + \int_{-\infty}^0 \langle y(s), y(s) \rangle ds = \|(y, h, u)\|^2 \end{aligned}$$

Hence ρ is a unitary representation of $\mathbb{R}$. Now note that from (7.2) we have that for the triple $(0, h, 0)$ the associated x is:

$$x(t) = \begin{cases} e^{itA_1}h, & t > 0 \\ e^{itA_1^*}h, & t < 0 \end{cases}.$$

Hence for positive t we obtain that $P\rho(t)(0, h, 0) = (0, x(t), 0) = (0, e^{itA_1}h, 0)$. $\square$

This idea leads us to consider the following construction. We construct weak solutions of the system of input and output compatibility equations from $\tilde{u}$ and $\tilde{y}$ and we plug these weak solution, more precisely the functions $\Lambda(\xi, \eta)(\tilde{u})$ and $\Lambda(\xi, \eta)(\tilde{y})$, into the formula (2.7) to get a state function x on $\mathbb{R}^d$ that is absolutely

continuous on lines $\xi t + \eta$, such that $\xi \in \text{Pos}(\mathfrak{V})$. However, first we need a dense subspace to work with.

Lemma 7.3. *Let $\mathcal{K}_0 \subset \mathcal{K}$ be the subspace of triples (y, h, u) , such that both $\tilde{y}$ and $\tilde{u}$ are twice continuously differentiable and both of the derivatives are square-summable, then $\overline{\mathcal{K}_0} = \mathcal{K}$.*

Proof. First note that since both $\tilde{u}$ and $\tilde{y}$ are twice continuously differentiable, we have that x is thrice continuously differentiable and $h = x(0)$. Using (7.2) we get that h is independent of u and y (it is in fact the initial condition). We must require that $\lim_{t \rightarrow 0+} u(t)$ and $\lim_{t \rightarrow 0-} y(t)$ exist and we denote them by $\tilde{u}(0)$ and $\tilde{y}(0)$, respectively. Similarly for their derivatives. Furthermore, we have the following condition on the values at 0 and the derivatives:

$$\tilde{u}(0) - \tilde{y}(0) = i\Phi h, \quad \tilde{u}'(0) - \tilde{y}'(0) = \Phi\Phi^*\tilde{u}(0) - \Phi A_1 h = \Phi\Phi^*\tilde{y}(0) - \Phi A_1^* h.$$

$$\tilde{u}''(0) - \tilde{y}''(0) = \Phi\Phi^*\sigma u'(0) - iA_1^2 h + iA_1\Phi^*\sigma u(0).$$

Therefore, a choice of $h \in \mathcal{H}$ forces three conditions on both u and y . However, a standard argument shows that twice continuously differentiable functions, with boundary conditions on them and their derivatives are dense in $L^2(\mathbb{R}_+, \mathcal{E})$ and $L^2(\mathbb{R}_-, \mathcal{E})$. $\square$

This lemma allows us to define for every $t \in \mathbb{R}^d$ an operator on $\mathcal{K}_0$. Let us denote by η_f the weak solution for the system of output compatibility equations with the initial condition f on the t_1 -axis and by $\Lambda_*(x, y)f$ the associated linear map. Given $(y, h, u) \in \mathcal{K}_0$, we construct $\tilde{u}$ and $\tilde{y}$. We then apply the transform u and η , respectively, to get (by Theorem 5.7) continuously differentiable functions $u^\dagger = u_{\tilde{u}}$ and $y^\dagger = \eta_{\tilde{u}}$ on $\mathbb{R}^d$, that solve the system of input and output compatibility conditions. Since these functions are continuously differentiable we can solve the associated system of our vessel with initial condition h , using (2.7) to obtain a twice continuously differentiable function x . Using the second equation of the system we obtain an output function z , that solves the system of output compatibility equations and coincides with $\tilde{y}$ on the t_1 -axis, thus by uniqueness $z = y^\dagger$. We define:

$$\rho(t)(y, h, u) = (y^\dagger(\cdot, t_2, \dots, t_d)|_{<t_1}, x(t), u^\dagger(\cdot, t_2, \dots, t_d)|_{>t_1}).$$

Note that it is immediate that $\rho(0)$ is the identity on $\mathcal{K}_0$.

Remark 7.4. Note that it is possible to use Equation (2.7) to construct the state signal for all $t \in \mathbb{R}^d$, since the operators $A_1, \dots, A_d$ are bounded and thus generate a group. In case these operators were unbounded one could run the adjoint system first to go back and then run the original system to obtain the value of the state signal. One of course would have in that case to show the commutation of the two actions.

Lemma 7.5. *For every $(y, h, u) \in \mathcal{K}_0$ we have that:*

- For every $t, s \in \mathbb{R}^d$, we have that $\rho(t)\rho(s)(y, h, u) = \rho(t+s)(y, h, u)$. Therefore, ρ is in fact a representation of $\mathbb{R}^d$.
- For every $t \in \text{Pos}(\mathfrak{V})$, we have that $\|\rho(t)(y, h, u)\| = \|(y, h, u)\|$ and hence, in particular $\rho(t)$ extends to an isometry on $\mathcal{K}$.
- For every $t \in -\text{Pos}(\mathfrak{V})$, we have that $\|\rho(t)(y, h, u)\| = \|(y, h, u)\|$ and hence $\rho(t)$ is in fact a unitary on $\mathcal{K}$, for every $t \in \text{Pos}(\mathfrak{V}) \cup -\text{Pos}(\mathfrak{V})$.

Proof. The first claim follows from the uniqueness part of Theorem 5.7. Namely, since $u^\dagger$ and $y^\dagger$ were determined uniquely by their restriction to any line with direction vector in $\text{Pos}(\mathfrak{V})$ and additionally restriction commutes with shifts, the claim follows.

For $t \in \mathbb{R}^d$ we write $\Lambda(t) = \Lambda(t, 0)$ and $\sigma(t) = \sum_{j=1}^d t_j \sigma_j$. To prove the second claim we apply (2.9) (modified to the straight line segment from 0 to t) to get that for every $t \in \text{Pos}(\mathfrak{V})$ we have:

$$\begin{aligned} \|x(t)\|^2 = \|h\|^2 + \int_0^1 \langle \sigma(t)(\Lambda(t)\tilde{u})(w), (\Lambda(t)\tilde{u})(w) \rangle dw - \\ \int_0^1 \langle \sigma(t)(\Lambda_*(t)\tilde{y})(w), (\Lambda_*(t)\tilde{y})(w) \rangle dw \end{aligned}$$

(note that by Theorem 5.7 we have that $\Lambda(t)(\tilde{u})(w) = u^\dagger(tw)$ and $\Lambda_*(t)(\tilde{y})(w) = y^\dagger(tw)$). Let us write $u_t = \Lambda(t)(\tilde{u})$ and $y_t = \Lambda_*(t)(\tilde{y})$. Applying Theorem 5.7 again we obtain:

$$\begin{aligned} \int_0^\infty \langle \sigma(t)u_t(w), u_t(w) \rangle dw &= \int_0^\infty \langle u(w), u(w) \rangle dw, \\ \int_{-\infty}^0 \langle \sigma(t)y_t(w), y_t(w) \rangle dw &= \int_{-\infty}^0 \langle y(w), y(w) \rangle dw. \end{aligned}$$

We now compute:

$$\begin{aligned} \|(y, h, u)\|^2 &= \|h\|^2 + \int_0^\infty \langle u(w), u(w) \rangle dw + \int_{-\infty}^0 \langle y(w), y(w) \rangle dw = \\ \|x(t)\|^2 - \int_0^1 \langle \sigma(t)u_t(w), u_t(w) \rangle dw + \int_0^1 \langle \sigma(t)y_t(w), y_t(w) \rangle dw + \\ \int_0^\infty \langle \sigma(t)u_t(w), u_t(w) \rangle dw + \int_{-\infty}^0 \langle \sigma(t)y_t(w), y_t(w) \rangle dw = \\ \|x(t)\|^2 + \int_1^\infty \langle \sigma(t)u_t(w), u_t(w) \rangle dw + \int_{-\infty}^{-1} \langle \sigma(t)y_t(w), y_t(w) \rangle dw = \\ \|x(t)\|^2 + \int_0^\infty \langle \sigma(t)(\Lambda(t, t)\tilde{u})(w), (\Lambda(t, t)\tilde{u})(w) \rangle dw + \\ \int_{-\infty}^0 \langle \sigma(t)(\Lambda_*(t, t)\tilde{y})(w), (\Lambda_*(t, t)\tilde{y})(w) \rangle dw. \end{aligned}$$

Another application of Theorem 5.7 gives us that both $\Lambda(t, t)\Lambda(e_1, t)^*$ and $\Lambda_*(t, t)\Lambda_*(e_1, t)^*$ are causal isometric isomorphisms and hence:

$$\|(y, h, u)\| = \|\rho(t)(y, h, u)\|.$$

The third claim is identical to the second but we exchange the roles of u and y . Since for $t \in \text{Pos}(\mathfrak{V}) \cup -\text{Pos}(\mathfrak{V})$ we have that $\rho(t)\rho(-t) = \rho(-t)\rho(t) = 1$ by the first part of the lemma, we conclude that $\rho(t)$ is a surjective isometry and hence a unitary and that $\rho(-t) = \rho(t)^*$. □

Lemma 7.6. *For every $(y, h, u) \in \mathcal{K}$, the following function is well defined for every $t \in \mathbb{R}^d$, $x(t) = P_{\mathcal{H}}\rho(t)(y, h, u)$. Furthermore, x is continuous and absolutely continuous*

on lines with a direction vector $\xi \in \text{Pos}(\mathfrak{V})$ and we have the equality:

$$(7.3) \quad x(\xi s + \eta) = e^{i(\xi s + \eta)A} \left(h + i \int_0^s e^{-i(\xi w + \eta)A} \Phi^* \sigma(\xi)(\Lambda(\xi) \tilde{u})(w) dw \right).$$

And thus for every line $\xi s + \eta$, with $\xi \in \text{Pos}(\mathfrak{V})$, we have:

$$\frac{dx}{ds} = i\xi Ax - i\Phi^* \sigma(\xi)(\Lambda(\xi, \eta) \tilde{u}).$$

Proof. For every $\epsilon > 0$ we choose $(y_0, h, u_0) \in \mathcal{K}_0$, such that $\|(y, h, u) - (y_0, h, u_0)\| < \epsilon$. Then for every $t \in \mathbb{R}^d$ we get $\|x_0(t) - P_{\mathcal{H}}\rho(t)(y, h, u)\| < \epsilon$, since $P_{\mathcal{H}}\rho(t)$ is a contraction. Since x_0 is continuous a standard $\epsilon/3$ argument shows that x is continuous.

If we prove that $x(\xi s + \eta)$ has the form described in (7.3), then we immediately see that x is absolutely continuous on those lines. Let us fix $\xi \in \text{Pos}(\mathfrak{V})$, then:

$$x_0(\xi s) = e^{i\xi s A} h + i \int_0^s e^{i(\xi - w\xi)A} \Phi^* \sigma(\xi)(\Lambda(\xi)(\tilde{u}_0)(w)) dw.$$

We note that for every $0 \leq w \leq 1$, we have $1 - w \geq 0$ and thus $e^{i(\xi - w\xi)A}$ is a contraction. Hence:

$$\begin{aligned} x_0(\xi s) - x(\xi s) &= \left\| \int_0^s e^{i(\xi - w\xi)A} \Phi^* \sigma(\xi)(\Lambda(\xi)(\tilde{u}_0 - \tilde{u})(w)) dw \right\| \leq \\ &C \int_0^1 \|\sigma(\xi)(\Lambda(\xi)(\tilde{u}_0 - \tilde{u})(w))\| dw \leq C \sqrt{\int_0^1 \|\sigma_t(\Lambda(\xi)(\tilde{u}_0 - \tilde{u})(w))\|^2 dw}. \end{aligned}$$

Now using functional calculus we note that there is a constant C' , such that for every $\xi \in \mathcal{E}$ the inequality $\|\sigma(t)\xi\|^2 \leq C' \langle \sigma(t)\xi, \xi \rangle$ holds. We thus conclude that:

$$\begin{aligned} &\left\| \int_0^s e^{i(\xi - w\xi)A} \Phi^* \sigma(\xi)(\Lambda(\xi)(\tilde{u}_0 - \tilde{u})(w)) dw \right\| \\ &\leq C \sqrt{C'} \sqrt{\int_0^\infty \langle \sigma(\xi)(\Lambda(\xi)(\tilde{u}_0 - \tilde{u})(w), (\Lambda(\xi)(\tilde{u}_0 - \tilde{u})(w)) \rangle dw} < C \sqrt{C'} \sqrt{\epsilon}. \end{aligned}$$

Letting ϵ tend to 0 we get the desired result. $\square$

Remark 7.7. For every line $\xi s + \eta$, with $\xi \in \text{Pos}(\mathfrak{V})$, we have that:

$$(\Lambda(\xi, \eta) \tilde{y})(s) = (\Lambda(\xi, \eta)(\tilde{u})(s) + i\Phi x(\xi s + \eta)).$$

Proof of Theorem 6.3. First we note that by Lemma 7.5 and the fact that $\text{Pos}(\mathfrak{V})$ spans $\mathbb{R}^d$ we get that ρ is a unitary representation of $\mathbb{R}^d$ on $\mathcal{K}$.

Now we need to check that if $P_{\mathcal{H}}: \mathcal{K} \rightarrow \mathcal{H}$ is the orthogonal projection, then $P_{\mathcal{H}}\rho(t)(0, h, 0) = e^{itA}h$, for every $h \in \mathcal{H}$. This, however, follows immediately from Lemma 7.6. $\square$

Lastly, we discuss the minimality of the unitary dilation that we have constructed. Recall that the dilation ρ is minimal if $\overline{\text{Span}\{\rho(t)\mathcal{H} \mid t \in \mathbb{R}^d\}} = \mathcal{K}$ or equivalently that there exists no ρ -invariant subspace of $\mathcal{H}^\perp$. For simplicity we shall assume that $A_1, \dots, A_d$ have the dissipative embedding property.

Lemma 7.8. Let $\mathcal{X}$ and $\mathcal{Y}$ be Hilbert spaces and $A: \mathcal{X} \rightarrow \mathcal{Y}$ be an injective bounded linear operator. Then the induced linear map $A: S'(\mathbb{R}^d, \mathcal{X}) \rightarrow S'(\mathbb{R}^d, \mathcal{Y})$ is injective.

Proof. We need to show that for every $\varphi \in \mathcal{S}'(\mathbb{R}^d, \mathcal{X})$, such that $A\varphi = 0$ we have that for every $f \in \mathcal{S}(\mathbb{R}^d, \mathcal{X})$, $\langle \varphi, f \rangle = 0$. Since $\mathcal{S}(\mathbb{R}^d, \mathcal{X}) \cong \mathcal{S}(\mathbb{R}^d) \hat{\otimes} \mathcal{X}$ if we show the claim for elementary tensors we will be done, since their span is dense. So given a function $f \in \mathcal{S}(\mathbb{R}^d)$ and $v \in \mathcal{X}$, we consider the function $f(t)v$. If $v \in \text{Im } A^*$, then $v = A^*w$ and we get:

$$\langle \varphi, f(t)v \rangle = \langle A\varphi, f(t)w \rangle = 0.$$

Since $\text{Im } A^*$ is dense, for every $v \in \mathcal{X}$, we can choose a sequence $v_n \in \text{Im } A^*$, that converges to v . Thus the sequence $f(t)v_n$ will converge in $\mathcal{S}(\mathbb{R}^d, \mathcal{X})$ to $f(t)v$. By continuity of φ we have that $\langle \varphi, f(t)v \rangle = 0$ for every $v \in \mathcal{X}$ and we are done. $\square$

Lemma 7.9. *The dilation obtained above is minimal if $\mathfrak{V}$ is weakly strict.*

Proof. Let us consider the subspace $\mathcal{L} = \overline{\text{Span}\{\rho(t)\mathcal{H} \mid t \in \mathbb{R}^d\}} \subset \mathcal{K}$ and assume the vessel $\mathfrak{V}$ is weakly strict. Consider the orthogonal complement $\mathcal{L}^\perp$ of $\mathcal{L}$. Since ρ is unitary and $\mathcal{L}$ is ρ invariant, we have that $\mathcal{L}^\perp$ is ρ -invariant as well. Note that every vector in $\mathcal{L}^\perp$ is of the form $(y, 0, u)$ and by invariance we have that the state function x we generate is identically 0. From Lemma 7.6 we get that for $\xi \in \text{Pos}(\mathfrak{V})$ $\Phi^*\sigma(\xi)u(t) = 0$ almost everywhere on every line in direction ξ . By Theorem 5.7 for every $\psi \in \mathcal{S}(\mathbb{R}^d, \mathcal{E})$ we have:

$$\langle \Phi^*\sigma(\xi)u_{\tilde{u}}, \psi \rangle = \int_{\xi^\perp} \int_{-\infty}^{\infty} \langle \Phi^*\sigma(\xi)(\Lambda(\xi, \eta)\tilde{u}(s), \psi(s\xi + \eta)) \rangle ds d\eta = 0.$$

Now we note that since $\mathfrak{V}$ is weakly strict we have that $\bigcap_{\xi \in \text{Pos}(\mathfrak{V})} \ker \Phi^*\sigma(\xi) = 0$. Thus if we consider the operator $(\Phi^*\sigma_1, \dots, \Phi^*\sigma_d): \mathcal{E} \rightarrow \mathcal{E}^d$, then by Lemma 7.8 it is injective and we conclude that $u_{\tilde{u}} = 0$. We need now only to deduce that $\tilde{u} = 0$. Let now $g \in \mathcal{S}(\mathbb{R}, \mathcal{E})$ and $h \in \mathcal{S}(\mathbb{R}^{d-1})$. We write $t' = (t_2, \dots, t_d)$ and define a function $f(t_1, \dots, t_d) = g(t_1)h(t')$ in $\mathcal{S}(\mathbb{R}^d, \mathcal{E})$ and we get:

$$0 = \langle u_{\tilde{u}}, f \rangle = \int_{\mathbb{R}^{d-1}} g(t') \langle L_{\tilde{u}}(t'), h \rangle dt'.$$

Now we assume that $\tilde{u} \neq 0$, then there exists $h \in \mathcal{S}(\mathbb{R}, \mathcal{E})$, such that $\langle \tilde{u}, h \rangle \neq 0$. Since $L_{\tilde{u}}$ is a smooth function there exists a $\delta > 0$, such that for every $t' \in \mathbb{R}^{d-1}$ if $\|t'\| < \delta$, then without loss of generality $\text{Re}\langle L_{\tilde{u}}(t'), h \rangle > 0$. Furthermore, we can choose $g(t')$ to be a bump function that is 1 on the ball of radius $\delta/2$ and is zero outside of a ball of radius δ and is always positive. Thus:

$$\text{Re} \int_{\mathbb{R}^{d-1}} g(t') \langle L_{\tilde{u}}(t'), h \rangle dt' = \int_{B_\delta} g(t') \text{Re}\langle L_{\tilde{u}}(t'), h \rangle dt' > 0.$$

This is a contradiction and thus $\tilde{u} = 0$. Since x we can conclude that $y = 0$ and we are done. $\square$

This condition is sufficient, but we can also get a necessary condition. To this end we need the following simple lemma:

Lemma 7.10. *Let $\mathcal{E}$ be a Hilbert space and $W \subset \mathcal{E}$ a closed subspace. Let α_j and β_j , $j = 1, \dots, d$ be operators on $\mathcal{E}$, such that for every $s \in \mathbb{C}$, and every $1 \leq j < k \leq d$ we have:*

$$[s\alpha_j + \beta_j, s\alpha_k + \beta_k] = 0.$$

Then the following are equivalent:

- (i) *For every $j = 1, \dots, d$ we have that $\alpha_j W \subset W$ and $\beta_j W \subset W$,*

- (ii) For every polynomial $p \in \mathbb{C}[z_1, \dots, z_d]$ and every $s \in \mathbb{C}$, we have that $p(s\alpha_1 + \beta_1, \dots, s\alpha_d + \beta_d)W \subset W$,
(iii) For every polynomial $p \in \mathbb{C}[z_1, \dots, z_d]$ and almost every $s \in \mathbb{C}$, we have that $p(s\alpha_1 + \beta_1, \dots, s\alpha_d + \beta_d)W \subset W$,

Proof. The equivalence of (i) and (ii) is obvious as well as the fact that (ii) implies (iii). To see that (iii) implies (ii) note that for every $\xi \in W$ and every polynomial $p \in \mathbb{C}[z_1, \dots, z_d]$ the following function is a continuous function in s :

$$s \mapsto p(s\alpha_1 + \beta_1, \dots, s\alpha_d + \beta_d)\xi.$$

Composing with the projection onto $\mathcal{E}/W$ we get a continuous function that is 0 almost everywhere and thus is identically 0. $\square$

So assume $\mathfrak{V}$ is not weakly strict and write $\mathcal{W} = \cap_{j=1}^d \ker \Phi^* \sigma_j$. Assume that there exists a vector $w \in \mathcal{W}$, such that for every $j = 1, \dots, d$ we have that $\alpha_j w \in \mathcal{W}$ and $\beta_j w \in W$. Fix some compact set $K = [a, b] \subset \mathbb{R}_{>0}$ and set $\tilde{u} = \mathcal{F}^{-1}(1_K w)$. Note that the values of u are all in $\mathcal{W}$ since $\mathcal{W}$ is closed. Furthermore, we have that:

$$u(t_1, \dots, t_d) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i \sum_{j=1}^d t_j (s\alpha_j + \beta_j)} \hat{u}(s) ds = \frac{1}{\sqrt{2\pi}} \int_a^b e^{i \sum_{j=1}^d t_j (s\alpha_j + \beta_j)} w ds.$$

Now the last expression belongs to $\mathcal{W}$ by our assumption. Then if we construct the associated state x for an initial condition h we get that $x = 0$ identically. Therefore, the triple $(u|_{\mathbb{R}_{<0}}, 0, u|_{\mathbb{R}_{>0}})$ is a non-zero vector orthogonal to $\mathcal{L}$, the space defined in the Lemma above.

Proposition 7.11. *If there exists a closed subspace $0 \neq \mathcal{M} \subset \mathcal{W}$, invariant under α_j and β_j for every $j = 1, \dots, d$, then the construction yields a non-minimal dilation. In other words if we have a minimal dilation then there is no such subspace of $\mathcal{W}$.*

REFERENCES

- [1] V. M. Adamjan and D. Z. Arov. On unitary coupling of semiunitary operators. *Amer. Math. Soc. Transl. (2)*, 13:75–129, 1970. English translation of the Russian original, *Mat. Issled.* 1 (1966) vyp. 2, 3-64.
- [2] Herbert Amann. *Linear and quasilinear parabolic problems, vol I*. Birkhäuser Verlag, Basel, 1995.
- [3] T. Andô. On a pair of commutative contractions. *Acta Sci. Math. (Szeged)*, 24:88–90, 1963.
- [4] T. Andô. Unitary dilation for a triple of commuting contractions. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 24(10):851–853, 1976.
- [5] J. Robert Archer. Positivity and the existence of unitary dilations of commuting contractions. In *The extended field of operator theory*, volume 171 of *Oper. Theory Adv. Appl.*, pages 17–35. Birkhäuser, Basel, 2007.
- [6] William Arveson. Subalgebras of C^* -algebras. II. *Acta Math.*, 128(3-4):271–308, 1972.
- [7] William Arveson. Dilation theory yesterday and today. In *A glimpse at Hilbert space operators*, volume 207 of *Oper. Theory Adv. Appl.*, pages 99–123. Birkhäuser Verlag, Basel, 2010.
- [8] William B. Arveson. Subalgebras of C^* -algebras. *Acta Math.*, 123:141–224, 1969.
- [9] Joseph A. Ball and Nir Cohen. de Branges-Rovnyak operator models and systems theory: a survey. In *Topics in matrix and operator theory (Rotterdam, 1989)*, volume 50 of *Oper. Theory Adv. Appl.*, pages 93–136. Birkhäuser, Basel, 1991.
- [10] Joseph A. Ball, Cora Sadosky, and Victor Vinnikov. Conservative linear systems, unitary colligations and Lax-Phillips scattering: multidimensional generalizations. *Internat. J. Control*, 77(9):802–811, 2004.
- [11] Joseph A. Ball and Victor Vinnikov. Overdetermined multidimensional systems: state space and frequency domain methods. In *Mathematical systems theory in biology, communications, computation, and finance (Notre Dame, IN, 2002)*, volume 134 of *IMA Vol. Math. Appl.*, pages 63–119. Springer, New York, 2003.

- [12] S. Brehmer. Über vetauschbare Kontraktionen des Hilbertschen Raumes. *Acta Sci. Math. Szeged*, 22:106–111, 1961.
- [13] M. S. Brodskii and M. S. Livšic. Spectral analysis of non-selfadjoint operators and intermediate systems. *Amer. Math. Soc. Transl. (2)*, 13:265–346, 1960. English translation of the Russian original, *Uspehi Mat. Nauk (N.S.)* 13 (1958) no. 1(79), 3–85.
- [14] Lawrence C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [15] Gerald B. Folland. *Introduction to partial differential equations*. Princeton University Press, Princeton, NJ, second edition, 1995.
- [16] Dumitru Gaşpar and Nicolae Suciu. On the intertwining of regular dilations. *Ann. Polon. Math.*, 66:105–121, 1997. Volume dedicated to the memory of Włodzimierz Mlak.
- [17] Dumitru Gaşpar and Nicolae Suciu. On the generalized von Neumann inequality. In *Recent advances in operator theory and related topics (Szeged, 1999)*, volume 127 of *Oper. Theory Adv. Appl.*, pages 291–304. Birkhäuser, Basel, 2001.
- [18] Anatolii Grinshpan, Dmitry S. Kaliuzhnyi-Verbovetskyi, Victor Vinnikov, and Hugo J. Woerdeman. Classes of tuples of commuting contractions satisfying the multivariable von Neumann inequality. *J. Funct. Anal.*, 256(9):3035–3054, 2009.
- [19] A. Grothendieck. *Topological vector spaces*. Gordon and Breach Science Publishers, New York-London-Paris, 1973. Translated from the French by Orlando Chaljub, Notes on Mathematics and its Applications.
- [20] Alexandre Grothendieck. Sur certains espaces de fonctions holomorphes. I and II. *J. Reine Angew. Math.*, 192:35–64, 77–95, 1953.
- [21] Paul R. Halmos. Normal dilations and extensions of operators. *Summa Brasil. Math.*, 2:125–134, 1950.
- [22] J. William Helton. Discrete time systems, operator models, and scattering theory. *J. Functional Analysis*, 16:15–38, 1974.
- [23] J. William Helton, Joseph A. Ball, Charles R. Johnson, and John N. Palmer. *Operator theory, analytic functions, matrices, and electrical engineering*, volume 68 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1987.
- [24] Peter D. Lax and Ralph S. Phillips. *Scattering theory*, volume 26 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, second edition, 1989. With appendices by Cathleen S. Morawetz and Georg Schmidt.
- [25] M. S. Livšic. Commuting nonselfadjoint operators and solutions of systems of partial differential equations generated by them. *Soobshch. Akad. Nauk Gruz. SSR*, 91(2):281–284, 1978.
- [26] M. S. Livšic. Operator waves in Hilbert space and related partial differential equations. *Integral Equations Operator Theory*, 2(1):25–47, 1979.
- [27] M. S. Livšic, N. Kravitsky, A. S. Markus, and V. Vinnikov. *Theory of commuting nonselfadjoint operators*, volume 332 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1995.
- [28] M. S. Livšic and L. L. Waksman. Open geometry, and operator colligations. *Ukrain. Geometr. Sb.*, (15):16–35, i, 1974.
- [29] Stephen Parrott. Unitary dilations for commuting contractions. *Pacific J. Math.*, 34:481–490, 1970.
- [30] B. Pavlov. Irreversibility, Lax-Phillips approach to resonance scattering and spectral analysis of non-self-adjoint operators in Hilbert space. *Internat. J. Theoret. Phys.*, 38(1):21–45, 1999. Irreversibility and cosmology. Fundamental aspects of quantum mechanics (Peyresq, 1997).
- [31] B. S. Pavlov. Selfadjoint dilation of a dissipative Schrödinger operator, and expansion in its eigenfunction. *Mat. Sb. (N.S.)*, 102(144)(4):511–536, 631, 1977.
- [32] Marek Ptak. Unitary dilations of multiparameter semigroups of operators. *Ann. Polon. Math.*, 45(3):237–243, 1985.
- [33] Donald Sarason. On spectral sets having connected complement. *Acta Sci. Math. (Szeged)*, 26:289–299, 1965.
- [34] J. J. Schäffer. On unitary dilations of contractions. *Proc. Amer. Math. Soc.*, 6:322, 1955.
- [35] Orr M. Shalit. Dilation theorems for contractive semigroups. *arXiv*, 1004.0723, 2010.
- [36] Eli Shamovich and Victor Vinnikov. Lie algebra operator vessels. *arXiv*, 1209.4224, 2014.
- [37] Eli Shamovich and Victor Vinnikov. Livsic-type determinantal representations and hyperbolicity. *arXiv*, 1410.2826, 2014.

- [38] M. Słociński. Unitary dilation of two-parameter semi-groups of contractions. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 22:1011–1014, 1974.
- [39] Marek Słociński. Unitary dilation of two-parameter semigroups of contractions. II. *Zeszyty Nauk. Uniw. Jagielloń. Prace Mat.*, (23):191–194, 1982.
- [40] Béla Sz.-Nagy. Sur les contractions de l'espace de Hilbert. *Acta Sci. Math. Szeged*, 15:87–92, 1953.
- [41] Béla Sz.-Nagy, Ciprian Foias, Hari Bercovici, and László Kérchy. *Harmonic analysis of operators on Hilbert space*. Universitext. Springer, New York, second edition, 2010.
- [42] François Trèves. *Topological vector spaces, distributions and kernels*. Academic Press, New York-London, 1967.
- [43] N. Th. Varopoulos. On an inequality of von Neumann and an application of the metric theory of tensor products to operators theory. *J. Functional Analysis*, 16:83–100, 1974.
- [44] Garth Warner. *Harmonic analysis on semi-simple Lie groups. I*. Springer-Verlag, New York, 1972. Die Grundlehren der mathematischen Wissenschaften, Band 188.

DEPARTMENT OF MATHEMATICS, TECHNION - ISRAEL INSTITUTE OF MATHEMATICS, HAIFA, 3200003, ISRAEL

E-mail address: shamovich@tx.technion.ac.il

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, 84105 BEER-SHEVA, ISRAEL

E-mail address: vinnikov@math.bgu.ac.il